Grand Valley State University ScholarWorks@GVSU

Masters Theses

Graduate Research and Creative Practice

6-2009

Fish Assemblages in Manistee River Tributaries: Longitudinal Distribution Analysis, Seasonal Variation, and Riparian Improvement Evaluation

Nicholas J. Gressick Grand Valley State University

Follow this and additional works at: http://scholarworks.gvsu.edu/theses



Over the Adviculture and Fisheries Commons, and the Biology Commons

Recommended Citation

Gressick, Nicholas J., "Fish Assemblages in Manistee River Tributaries: Longitudinal Distribution Analysis, Seasonal Variation, and Riparian Improvement Evaluation" (2009). Masters Theses. Paper 3.

This Thesis is brought to you for free and open access by the Graduate Research and Creative Practice at ScholarWorks@GVSU. It has been accepted for inclusion in Masters Theses by an authorized administrator of ScholarWorks@GVSU. For more information, please contact scholarworks@gvsu.edu.

FISH ASSEMBLAGES IN MANISTEE RIVER TRIBUTARIES: LONGITUDINAL DISTRIBUTION ANALYSIS, SEASONAL VARIATION, AND RIPARIAN IMPROVEMENT EVALUATION

A thesis submitted in partial fulfillment of the requirements for the degree of Master of Science

Ву

Nicholas J. Gressick

To


Biology Department Grand Valley State University Allendale, Michigan June 2009 Signature page has been removed.



Copyright by

Nicholas John Gressick

2009



ACKNOWLEDGEMENTS

I wish to thank Dr. Eric Snyder, Dr. Mark Luttenton, Dr. Carl Ruetz, and Marty Holtgren for their support and guidance. I would also like to thank Grand Valley State University for this great opportunity. Also, the Little River Band of Ottawa Indians for assistance in the field and use of equipment. I would like to thank Kevin Donner, April Wright, Stephanie Ogren, and Bob Sanders for assistance with sampling. I would also like to thank the Environmental Protection Agency for providing funds available for this project. Also, I thank the Little River Band of Ottawa Indians and Grand Valley State University for equipment and guidance with the project.

Abstract

Sedimentation and culvert issues can affect both stream physical and biological integrity and can negatively impact the fish assemblage by impeding fish passage, degrading food resource availability, and masking requisite spawning gravel and cobble. The purpose of this study included (1) an attempt to quantify impacts of poorly constructed road stream crossings and eroding banks on fish assemblages and subsequently assess these sites as sediment sources and connectivity breaks on entire fish assemblages and individual fish species, (2) give a detailed description of the fish assemblage structure from headwaters to mouth on the three study streams and assess the differences between streams regarding fish assemblage structure, and (3) designate the potential differences between whole fish assemblages and individual species between three tributaries of the lower Manistee River, Michigan, using a headwater to mouth approach. Electrofishing was conducted during spring and fall 2004 and 2005 on three tributary streams (Sickle Creek, 1st order, Pine Creek, 2nd order, Bear Creek, 4th order) within the lower Manistee River watershed. A total of 29 electrofishing reaches were sampled and included 5 road-stream and streambank restoration sites. Sickle Creek had reduced diversity and increased dominance above a substantially perched culvert. Fish assemblage response above and below impact sites was mixed, and largely determined by either an up vs. downstream impact. In the longitudinal analysis, unique fish assemblages were observed between Sickle, Pine, and Bear Creeks. Whole fish assemblage measurements revealed no significant differences between seasons for Pine, Sickle, or Bear Creeks with respect to fish density, dominance, diversity, and richness. It appears that different order streams and patterns in fish community abundance and diversity seemed to reflect the environmental habitat template, even when this template deviated from the predicted longitudinal conditions.

TABLE OF CONTENTS

LIST	OF TABLES	xi		
LIST	LIST OF FIGURES x			
СНА	PTER			
I.	Preface	1		
II.	Response of fish assemblage structure to historic sand in	nputs and		
	consequences for restoration	5		
	a. Abstract	5		
	b. Introduction	5		
	c. Materials/methods	8		
	i. Study area	8		
	ii. Study stream characteristics	8		
	iii. Experimental design	10		
	d. Results and Discussion	12		
	i. Whole fish assemblage impacts	12		
	ii. Taxa-specific patterns	15		
	e. Conclusion	21		
	f. Literature cited	22		
	g. Tables and figures	28		
III.	Longitudinal fish distribution in three West Michigan streams of different			
	orders	40		
	a. Abstract	40		
	b. Introduction	40		
	c. Materials/methods	43		
	i. Experimental design	43		
	d. Results and Discussion	44		
	i. Longitudinal predictions	44		
	ii. Fish assemblages and water chemistry	49		
	e. Conclusion	51		
	f. Literature cited	52		

	g. Tables and figures	54
IV.	Fish assemblage spring/summer longitudinal patterns in three tribu	itaries of
	the lower Manistee River, Michigan	69
	a. Abstract	69
	b. Introduction	69
	c. Materials/methods	71
	i. Experimental design	71
	d. Results and Discussion	72
	i. Whole fish community seasonal differences	72
	ii. Taxa-specific seasonal patterns	72
	e. Conclusion	77
	f. Literature cited	78
	g. Tables and figures	81
V.	Overall discussion	89
APPE	NDICES	
	a. Appendix A – Electrofishing data	93
	LIST OF TABLES	
Table	1. Mean water quality variables for each respective reach sampled.	28
	2. Average abundance data for most influential species affecting site ences in NMDS.	e
		28
	3. Mean density, CPUE, and relative abundance data for significant ences between upstream and downstream locations at respective site	
	4. Summary of various taxa-specific and community level responses and below restoration sites.	s 29
Table	5. Set-up and results of NMDS analysis for each restoration site.	30
	6. Analysis of variance results for various species CPUE for udinal distribution in Sickle Creek.	54
	7. Analysis of variance results for various species CPUE for udinal distribution in Pine Creek.	54

Table 8. Analysis of variance results for various species CPUE for longitudinal distribution in Bear Creek.	55
Table 9. Fish species that differed significantly between upper, mid, and lower reaches for Sickle, Pine, and Bear Creeks.	55
Table 10. Data values for significant differences between seasons.	81
Table 11. Summary of spring/summer fish community responses in Sickle, Pine, and Bear Creeks.	81
LIST OF FIGURES	
Figure 1. Sickle, Pine, and Bear Creeks within the Manistee River Watershed.	31
Figure 2. Sickle Creek site design.	31
Figure 3. Total fish density above and below road stream crossings.	32
Figure 4. Fish species dominance above and below road stream crossings.	33
Figure 5. Species diversity above and below road stream crossings.	34
Figure 6. Species richness above and below road stream crossings.	35
Figure 7. Fish assemblage differences among Pine Lake Road reaches, Pine Creek, based on NMDS analysis.	36
Figure 8. Fish assemblage differences among Steinberg Road reaches, Pine Creek.	37
Figure 9. Fish assemblage differences at Sickle Creek.	38
Figure 10. Fish assemblage differences among Milks Road reaches, Bear Creek.	39
Figure 11. Sickle Creek longitudinal profile for Spring 2004, Fall 2004, Spring 2005, and Fall 2005.	56
Figure 12. Pine Creek longitudinal profile for Spring 2004, Fall 2004, Spring 2005, and Fall 2005.	57

Figure 13. Bear Creek longitudinal profile for Fall 2004, Spring 2005, and Fall 2005.	58
Figure 14. Catch per unit effort data for various fish species in Sickle Creek.	59
Figure 15. Catch per unit effort data for various species in Pine Creek.	60
Figure 16. Catch per unit effort data for various fish species in Bear Creek.	61
Figure 17. Correlation between discharge and species richness and fish density in Sickle Creek.	62
Figure 18. Discharge was negatively correlated with species richness and species diversity in Pine Creek.	63
Figure 19. Temperature was negatively correlated with species dominance and species diversity in Bear Creek. Dissolved oxygen was also negatively correlated with species diversity.	
Figure 20. NMDS plot showing dispersal of fish assemblages at reaches and their relationship to water quality measurements in Sickle Creek.	66
Figure 21. NMDS plot showing dispersal of fish assemblages of reaches and their relationship towards water quality measurements in Pine Creek.	67
Figure 22. NMDS plot showing dispersal of fish assemblages of reaches and their relationship towards water quality measurements in Bear Creek.	68
Figure 23. Relative seasonal fish density of Pine, Sickle, and Bear Creeks, Manistee County, Michigan.	82
Figure 24. Seasonal fish dominance for Pine, Sickle, and Bear Creeks, Manistee County, Michigan.	83
Figure 25. Seasonal species diversity of Pine, Sickle, and Bear Creeks, Manistee County, Michigan.	84
Figure 26. Seasonal fish species richness of Pine, Sickle, and Bear Creeks, Manistee County, Michigan.	85
Figure 27. Relative fish density per hectare of all significant fish differences for each respective stream.	86

Figure 28. Fish catch per minute of all significant fish differences for each respective stream.	1 87
Figure 29. Fish relative abundance of all significant fish differences for Bear Creek.	88

CHAPTER 1

PREFACE

In 2003, the Little River Band of Ottawa Indians received a U.S. Environmental Protection Agency Targeted Watershed Initiative grant to aggressively improve water quality in the Big Manistee River watershed. Restoration efforts focused on the Bear Creek and Pine Creek sub-watersheds. Projects included 1) replacing undersized culverts with the appropriate open bottom culvert or bridge and 2) replacing undersized bridges with modern bridges designed to direct stormwater (and sediment run-off) away from the stream. Additionally, these designs improved fish passage and restored the natural flow patterns to help remove accumulated fine sediment (silt and sand). The main objectives were to: 1) determine site locations for road-stream crossings, streambanks and accesssite improvements through review of pre-existing inventories or site assessments, 2) implement restoration activities of constructing new road-stream crossings, stabilizing streambanks and creating access-sites, and 3) to critically evaluate the effectiveness of each habitat restoration practice through the use and development of biological metrics. The focus of this thesis project revolved around the third objective and specifically involved an assessment of the fish assemblages while two other GVSU graduate students, Nichol De Mol and April Wright, focused on aquatic insect communities and sediment composition change, respectively.

Excessive sedimentation can be considered a major abiotic disturbance affecting stream biotic community structure and function (Lake 2000). In fact, sediment has been found to adversely affect most fish species, especially salmonids by smothering redds,

masking spawning gravel and reducing habitat complexity (Barton 1977, Scott et al. 1986, Alexander and Hansen 1988, Curry and MacNeil 2004, and many others).

Most of the studies previously mentioned used a BACI design, however, only used simple comparisons in relation to sedimentation problems. In this study we used an upstream versus downstream comparison, with multiple sampling points at both upstream and downstream sites. Fausch et al. (2002) suggested that the importance of different physical and ecological processes will be revealed at different spatiotemporal scales, and processes will interact among scales. Furthermore, Allan (2004) suggested that further research is needed that examines responses to land-use under different management strategies and that employs response variables that have greater diagnostic value than many of the current measures. The results of our study, in conjunction with the considerations in the previous two studies, contribute to our understanding of what occurs further upstream and downstream of a potential sediment source, although results of this research constitute the pre-restoration monitoring—additional GVSU MS theses document the post-restoration phase (De Mol 2007, DeBoer 2008).

In addition to a larger spatial assessment of sedimentation, we have utilized a headwaters to mouth approach within the Great Lakes Region to examine stream fish distribution impaired by high rates of sedimentation and barriers to fish migration. Many studies have examined fish longitudinal distribution and its relationship to biotic and abiotic factors (Schlosser 1991, Grenouillet et al. 2004, Schaefer and Kerfoot 2004, Helms et al. 2008 and many others). Schlosser (1991) stated that, as a whole, land-use activities can decrease spatial heterogeneity and connectivity of physical habitats. Torgersen et al. (2006) compared gradients in fish assemblage structure among rivers and

at multiple spatial scales and found spatial structuring of fish assemblages exhibited a generalized pattern of cold- and coolwater fish assemblage zones, but was variable between thermal zones, particularly in the warmest stream. Helms et al. (2008) evaluated the impact of land cover on fish assemblages in a western Georgia stream and found that fish assemblages were largely explained by physiochemical and hydrological rather than habitat variables. Similar to our study area, Zorn et al. (1998) examined the distribution and abundance patterns of fish assemblages at numerous locations in lower Michigan streams using low-flow yield and catchment area as variables. While providing a framework for stream fish distribution in the lower peninsula, they also determined that stream fishes respond in an individualistic manner to stream conditions, and that focus on individual species is needed to describe fish assemblage structure in streams. While each of these studies attempted to explain fish distribution in each stream using various abiotic measurements with few in sand-dominated, upper Midwest streams, we examined the relationship with Michigan coldwater stream fish longitudinal distribution and various water quality parameters, substrate composition, up vs. downstream impacts of localized sediment sources, and stream passage constrictions created by road stream crossings and eroding stream banks.

The purpose of this study was to: 1) examine the response of fish assemblage structure to historic sand inputs, constricted and perched culverts, and consequences for restoration, 2) study longitudinal fish distribution from up to downstream, and 3) focus on fish assemblage spring/summer longitudinal patterns in three tributaries of the lower Manistee River, Michigan.

This thesis is divided into three chapters that focus on sediment and restoration, longitudinal fish assemblage, and seasonal (spring and summer) fish assemblage. Finally, all three chapters are summarized.

CHAPTER 2

RESPONSE OF FISH ASSEMBLAGE STRUCTURE TO HISTORIC SAND INPUTS, CONSTRICTED AND PERCHED CULVERTS, AND CONSEQUENCES FOR RESTORATION

ABSTRACT

Sedimentation affects both stream physical and biological integrity. Improperly designed stream passage accompanied with sedimentation and altered hydrology can impede fish passage and reduce fish assemblage integrity. The purpose of this study was to: 1) quantify impacts of poorly constructed road stream crossings and eroding banks on fish assemblages, and 2) assess these sites as sediment sources and connectivity breaks on entire fish assemblages and individual fish species. Electrofishing was conducted during spring and fall 2004 and 2005. A total of 29 electrofishing reaches were sampled which included 5 road-stream and streambank restoration sites. Sickle Creek (1st order) had reduced diversity and increased dominance above a substantially perched culvert (Shannon's diversity = 0.180 vs. 0.552; Simpson's dominance = 0.688 and 0.412 above vs. below, respectively). Pine Creek (2nd order) had 12 reaches sampled, while Bear Creek (4th order) had 7 reaches sampled. In both river systems, fish assemblage response above and below impact sites was mixed, and largely determined by either an up vs. downstream impact. For example, undersized road-stream culverts reduced upstream habitat quality while eroding banks reduced downstream habitat quality. Improvements to road-stream crossings should be done to maximize natural river structure and function.

Introduction:

Stream disturbance plays a major role in determining the structure and function of stream communities (Lake 2000). Disturbance determines both patchiness and diversity in streams (Lake 2000). Sedimentation represents a disturbance that can negatively impact fish assemblages by altering physical, chemical, and biological characteristics of streams (Lake 2000).

Although many studies have examined the effects of natural and anthropogenic disturbance on streams, there are fewer studies involving fish assemblages and

sedimentation effects from road stream crossings, especially in a longitudinal gradient perspective. Whole watershed disturbances include water fluctuations, forestry practice, livestock grazing, and urbanization, all of which can contribute to sedimentation (Larimore et al. 1959, Junk et al. 1989, Jones et al. 1999). Local or proximate disturbances such as sedimentation can affect fish assemblages directly by changing water quality, habitat, and spawning areas (Bjornn 1971, Berg and Northcote 1985, Lisle 1989, Lisle and Hilton 1991, Servizi and Martens 1992, Waters 1995).

Peters (1967) found that brown trout in a Montana stream decreased with increased sedimentation from agricultural sources, similar to results obtained by Saunders and Smith (1965) who studied the effect of heavy siltation in a Prince Edward Island brook trout stream. They reported 70% declines in trout populations, (both age-0 and older fish) due to loss of cover by sediment deposits. More recently, Alexander and Hansen (1988) showed an experimental addition of sand in Hunt Creek, Michigan, reduced the brook trout population by 50%. Drastic declines in a Minnesota brook trout population after catastrophic spring floods were observed from stream bottom sediments covered by shifting sand, leading to scouring of eggs and reducing fry abundance (Elwood and Waters 1969). Curry and Macneill (2004) studied sediment deposition effects on brook trout embryos and young of the year. They found that mortality occurred at late encapsulated stages before hatching; the result of depleted oxygen from fine sediment deposition. Groundwater inputs in some areas reduced sedimentation in redds and increased survival (Curry and Macneill 2004)

Sedimentation can occur when road-stream crossings are constructed. For example, Beschta (1978) showed that road construction causes soil movement or

landslides. Chisholm and Downs (1978) reported that construction of a four-lane highway along Turtle Creek, West Virginia generated large amounts of sediment, burying the streambed under 10 inch deposits and eliminating the stream benthos. Road construction along Joe Wright Creek, a mountain stream in Colorado, reduced macroinvertebrates where sediment deposits occurred (Cline et al. 1983). Loomis (1989) showed detrimental impacts of road building on recreational and commercial fisheries. Barton (1977) observed a decline in total fish standing stock from 24 to 10 kg/ha owing to sedimentation downstream from a bridge construction on an Ontario brook trout stream. Comparative analysis of a pristine creek and a road-stream crossing impacted by urban development, suggested restructuring of the fish community in Kelsey Creek, Washington (Scott et al. 1986). Specifically, results showed environmental disturbance, including habitat alteration (sedimentation), increased nutrient loading, and degradation of the intragravel environment had a large negative effect on coho salmon, while not affecting cutthroat trout (Oncorhynchus clarki). Kelsey Creek became dominated with cutthroat trout, whereas the control stream had a diverse composition of salmonids.

In summary, research shows that species-specific responses to road-stream crossings are variable and urban development does not necessarily displace all salmonids. However, these studies did not account for overall stream ecosystem integrity when road-stream crossing construction occurred. Alternatively, if construction is conducted in an ecologically appropriate manner, then impacts can be potentially minimized. Improvement of existing road-stream crossings that historically contributed significant amounts of sediment could substantially improve the fish assemblage by reducing sediment impacts.

The purpose of this study was to: 1) quantify impacts of existing road-stream crossings and eroding banks on fish assemblages, and 2) assess these sites as sources of sediment and connectivity breaks on entire fish assemblages and individual fish species.

Materials and Methods:

Study Area: The Manistee River flows westward into Lake Michigan through northern lower Michigan (Figure 1) through the Manistee National Forest and drains a largely forested watershed in northwestern lower Michigan characterized by a stable baseflow. Currently, land use is mainly forested (75%) with occasional agricultural practice (25%). Sickle, Pine, and Bear creek tributaries of the lower Manistee River (Figure 1) were chosen based on their unique fish assemblages and accessibility.

Study Stream characteristics: Sickle Creek is a small 1st order stream that flows southward into the Big Manistee River. Discharge in this stream can fluctuate rapidly and average wetted stream width is 2 m. Riparian vegetation consists mainly of white cedar (Thuja occidentalis), American basswood (Tilia americana), silver maple (Acer saccharinum), red maple (Acer rubrum), and musclewood (Carpinus caroliniana). Canopy cover is approximately 80% with associated large woody debris and sediment composition consists of the largest component of sand, with occasional silt, pebble and gravel pockets. The fish community includes brook trout, brown trout, and rainbow trout, along with juvenile Chinook salmon (Oncorhynchus tshawytscha) and coho salmon as well as burbot and associated resident fish species including mottled sculpin (Cottus bairdi). In each stream, dissolved oxygen (mg/L), pH, specific conductivity (umhos/m), and temperature (OC) were measured using portable water quality multimeters (either a

Hydrolab and/or YSI 600). Discharge was measured using a Marsh-McBirney flow meter (Table 1).

Pine Creek is a moderately-sized, 2nd order stream that originates at Pine Lake and flows northwest to the Manistee River. Discharge fluctuates seasonally with stream widths averaging 5m. Riparian vegetation at lower regions includes white cedar, maples, eastern hemlock (*Tsuga canadensis*) and American basswood with upper regions primarily consisting of speckled alder (*Alnus rugosa*), white pine (*Pinus strobus*), white cedar, red osier dogwood (*Cornus stolonifera*) and silky dogwood (*Cornus amomum*). Canopy cover is approximately 50%, and large woody debris is abundant. Sediment composition consists largely of sand, with additional silt, pebble, and gravel. Pine Creek has a self-sustaining brown trout population and resident fish species including mottled sculpin, rainbow trout, and western blacknose dace.

Bear Creek is a large, 4th order stream that flows southward to the Manistee River. Discharge fluctuates seasonally with a 10 m average wetted stream width. Riparian vegetation at lower regions include maples, American basswood, willow (*Salix spp.*), and sassafras (*Sassafras albidum*). Stream riparian vegetation includes white cedar, speckled alder, and American basswood. Canopy cover is higher (75%) at the lower regions than at upper regions (50%). Large woody debris is moderate in lower and upper regions and stream sediments composition consists of mainly sand, along with some silt, while upper regions consist of sand, silt, pebble, gravel, and cobble. Bear Creek has self-sustaining brown trout and brook trout populations, and resident fish species include longnose dace and mottled sculpin.

Experimental design: Study reaches were delineated within each stream; 5 in Sickle, 7 in Bear, and 12 in Pine creeks. The number of reaches was determined based on stream size, length, and accessibility. The basic sampling design for Sickle Creek (Figure 2) is also representative of Pine and Bear creeks. Each reach within each stream represented a replicate, and each of the replicates were used in the statistical analyses with 2 replicates for each reach for each respective season, with a total of 4 per reach for the two years of the study, excluding the Bear Creek reaches that lacked a spring 2004 sampling period due to high water and equipment problems leaving one replicate for spring and 2 for fall.

Electrofishing reach length was determined using a value of 40 times stream width, which is the standard used by the EPA (Barbour et al. 1999). Fish assemblages were assessed using pulsed DC electrofishing equipment (Carline 2001). A Smith-Root backpack electrofisher (Model 15-D 300 Watt generator-powered) was used to sample Sickle and Pine Creeks, while Bear Creek was sampled with a tote barge (Model 15-D 300 Watt generator-powered). Sickle Creek, the smallest system, was sampled using a multiple-pass depletion technique with blocker nets, while the two larger systems, Pine Creek (2nd order) and Bear Creek (4th order), were sampled using single-pass techniques with no blocker nets. Allen at al. (2003) showed that the single-pass technique was as effective as the multiple-pass technique to sample fish assemblages.

Whole Fish Assemblages and Individual Species: Fish assemblages were estimated for each respective stream during spring and summer 2004 and 2005 with the exception of Bear Creek in spring 2004 due to a combination of equipment problems and high water levels. Data were quantified using catch per unit effort (CPUE, i.e. fish/minute), which

divides catch by effort, removing the effect of variable effort in the abundance measurements (Hubert and Kohler 1999). However, it is important to note that CPUE may be proportional to abundance, but can be influenced by catchability of a species. Catchability may be influenced by gear selectivity, size and age of fish, horizontal and vertical distribution of fish, and electrofishing method (Maunder at al. 2006).

Total fish density, dominance, diversity, and richness were evaluated using repeated measures ANOVA (SAS 9.1) for whole fish assemblages using upstream vs. downstream fish assemblage data. Individual fish species were assessed using density, CPUE, and relative abundance. Significance was determined at p≤0.05. Density was calculated as the total number of the entire fish assemblage and individual species divided by the reach area although population estimates were not quantified. Dominance values were calculated using Simpson's Dominance Index, while diversity was calculated using Shannon's Diversity Index (Lake 2000). Richness was calculated as the sum of all species present at each reach.

Non-metric multidimensional Scaling (NMDS) was used to determine if there was any separation between fish communities at the four road-stream crossings (PCORD software package) (Mather 1976, Kruskal 1964). NMDS is an ordination method well suited to data that are nonnormal or on arbitrary, discontinuous, or otherwise questionable scales (McCune and Grace 2008). NMDS allows the user to avoid the assumption of linear relationships among variables, uses ranked distances to linearize the relationship between distances measured in species space and distances in environmental space, and allows the use of any distance measure. Historically, NMDS tended to fail to find the best solution because of intervening local minima, however, this problem is easily avoided by

requesting more random starts, more iterations, and a more stable solution. Scree plots were generated to determine the appropriate number of dimensions that should be used with each analysis and the main matrix used in NMDS was relative abundance.

Results and Discussion:

Whole Fish Assemblages: Pine Creek at Pine Lake Road showed significantly higher dominance values for downstream sites than upstream sites (Table 3). Differences were also seen in general fish assemblage structure between upstream and downstream reaches (Figure 3 and 7). These differences were based primarily on brook stickleback (*Culaea inconstans*), rainbow trout, and white sucker (*Catostomus commersoni*). Brook stickleback appeared to have higher abundance upstream with more suitable habitat. White sucker also had higher abundance upstream likely the result of upstream migration of juveniles. Rainbow trout were higher downstream where there was a larger abundance woody debris (personal observation) and larger sized substrate (DeMol 2007).

Possible reasons for the differences above and below the culvert based on qualitative observations include different habitat types, less fishing pressure up- vs. down-stream, and more LWD downstream. In addition, substrate was quantitatively measured and was significantly coarser downstream (A. Wright, personal communication, DeMol 2007). No significant differences were seen with total fish density or richness at this site.

Pine Creek at Steinberg Road showed no significant differences with respect to total fish density, dominance, diversity, and richness (CPUE, Table 3, Figures 4-7). However, study reaches were distinct with respect to the overall fish community (Figure

8). These differences were seen mainly with brown trout, mottled sculpin, and rainbow trout with higher abundance and CPUE downstream (Table 2). Qualitative differences suggested that there was more overhanging cover, while quantitative differences indicated significantly larger substrate downstream (A. Wright, personal communication, DeMol 2007). Grenouillet et al. (2004) showed that only stream width and gradient influenced local species richness and also that the relative importance of local habitat and biotic processes may depend on the position along the longitudinal gradient.

Other qualitative observations suggest that passage to upstream reaches at this location consisted of a constricted area with flow through non-even culverts. Depth of water within culverts was less than that directly downstream and upstream of the road crossing with considerable sediment buildup directly upstream with water backed up into a drainage ditch. Other differences could result from individual species requirements of substrate, habitat, water quality, food availability, and discharge. Directly downstream, larger substrate was evident, indicating passage of sediment further downstream from the road crossing. This indicates that poor quality substrate is not always evident below poorly constructed road stream crossings and increased water velocities are a likely cause of scouring directly downstream of the culverts.

The Sickle Creek site showed significantly greater density, diversity, and richness (CPUE) with downstream sites compared to upstream sites. Dominance was greater at the upstream locations, due to a higher relative abundance of mottled sculpin. Differences in general fish assemblage structure were also seen (Figure 9). Here, the two upstream reaches showed different fish assemblage structure compared to that of the downstream reaches. These differences were mostly attributed to increased abundance of burbot,

Chinook and coho salmon (CPUE) downstream (Table 2). Burbot are not capable of swimming against high velocities, which would have been the only method for breaching the severely perched culvert at this location (Wootton 1998). The Chinook and coho salmon were primarily young individuals and were likely not capable of swimming against the highest velocities that were present in spring. Additionally, during base flow, the culvert was perched by a large enough gap (approximately 3-4 inches) that these species could likely not pass upstream. Sickle Creek also empties directly into the Big Manistee River, providing adequate passage for species migrating in and out for refuge purposes, while other species are continuously in the creek.

Bear Creek at Milks Road showed no differences up- vs. down-stream for density, dominance, diversity, richness (CPUE). Results of the NMDS analysis suggested that the upstream Milks Road reach and the two downstream reaches were different from the first downstream reach (MILKSDS, Figure 10). These differences could be attributed to specific taxa including brown trout, mottled sculpin, and rainbow trout abundance which were higher downstream (Table 2). This road crossing also was undersized, but not to the same extent as the other road crossings and in addition there was a significant eroding bank just upstream of the culvert that was stabilized in the summer of 2007. Consequently, these species were able to move upstream to reach potentially more suitable habitat and better water quality. Further downstream from the Milks Road site, streambank stabilization measures were occurring on two sites, SWAINDS1 and SWAINDS2, which may have influenced fish assemblage structure further upstream at the Milks Road stream crossing.

Fish assemblage measurements in response to poorly constructed road crossings can be affected by fine sediment deposition as many species are affected by this size of sediment. Lisle and Hilton (1991) showed that filling of fines affects pool habitat by reducing volume, particularly during drought conditions, and covers substrate. Stream velocity can be linked to bed mobility and sediment transport and thereby to spawning habitat through scour of spawning substrate. Furthermore, fine sediment in pools is transported first as flow increases and could infiltrate spawning habitat constructed immediately downstream and adjacent riffles.

My results indicate that 3 of the 4 restoration sites included in this analysis experienced significant sediment impacts upstream—likely a result of undersized and poorly aligned culverts. These included the two sites on Pine Creek (Pine Lake Road and Steinberg) and the site on Sickle Creek. In contrast, the site on Bear Creek (Milks Road) did not show dramatic differences above vs. below an undersized bridge. Ongoing analyses beyond the scope of this thesis project have documented the rapid redistribution of fine sediment downstream once the culverts were replaced. Given enough time, the return of a more natural flow regime should see an improvement both up and downstream of these culverts as the system continues to establish a new dynamic equilibrium between flow and sediment transport.

Taxa-specific Patterns:

Individual Species at Pine Lake Road, Pine Creek: Western blacknose dace, mottled sculpin, and rainbow trout had significantly higher density, CPUE, and relative proportions at the downstream sites than the upstream sites (Table 3, 4). Higher proportions of large woody debris, coarse substrate, more suitable water quality

parameters, velocity, and food availability are likely the cause. Lisle and Hilton (1991) found that fine sediment deposition reduced fish pool habitat and obliterated substrate cover of certain fish species.

Brook stickleback, central mudminnow (*Umbra limi*), and white sucker were significantly higher in the upstream sites than the downstream sites in all measurements. Some possible factors include less competition for food, better species-specific water quality needs, lower velocities suitable for these species, and finer substrate. Creek Chub (*Semotilus atromaculatus*) relative proportions were significantly higher in the upstream sites but density and CPUE were not significant. Lower proportions of other species are possible contributing factors. Brook trout, brown trout, coho salmon, johnny darter (*Etheostoma nigrum*), longnose dace, and northern redbelly dace (*Phoxinus eos*) were not different in any of the measurements. Burbot, Chinook salmon, and northern brook lamprey (*Ichthyomyzon fossor*) were absent from this site possibly as a result of substrate, discharge, water quality, access to site, competition, and food availability.

Individual Species at Steinberg Road, Pine Creek: Western blacknose dace had significantly higher relative proportions upstream than downstream, although no significant differences were seen regarding density and CPUE (Table 3, 4). Western blacknose dace contributed a higher proportion of the upstream fish assemblage. Brown trout and rainbow trout density and relative proportion were significantly higher downstream, but CPUE was not. These species were better adapted for faster velocities and resided in conditions with larger amounts of substrate, woody debris, undercut banks, and possibly optimal food availability. McRae and Diana (2005) indicated that percent gravel substrate and percent emergent vegetation accounted for 62% of the variance in

age-0 brown trout density in the Au Sable River, Michigan. Waters (1992) indicated that stream salmonid production is influenced by water quality. But controversially, Mann and Penczak (1986) suggested that the potential level of salmonid production predicted is not by a stream's water quality, ultimately showing that more salmonids inhabit streams of lower water quality.

Substrate is very important to survival of some species. Bjornn (1971) showed that reductions in salmonid fry were linearly related to the degree of cobble embeddedness. The greatest effect of excess sediment occurred in pools, where decreases in area and depth caused decreases in summer rearing capacity for juvenile salmonids. Slower velocities at the Steinberg Road site influenced sediment-holding capacity in pools above the culverts.

Brook stickleback, brook trout, burbot, central mudminnow, Chinook salmon, coho salmon, creek chub, johnny darter, mottled sculpin, northern redbelly dace, and white sucker had no significant differences among all measurements. Even though differences in brook trout were not significant in any of the measurements, other studies suggest that mean minimum water temperatures and mean daily water temperature fluctuation account for variances in density of age-0 brook trout (McRae and Diana 2005). Longnose dace and northern brook lamprey were absent at this site possibly as a result of discharge, habitat, substrate, physiological components, water quality, and food availability.

Individual Species at USFS 5575, Sickle Creek: Brook trout relative proportions were significantly higher at the upstream sites versus the downstream but density and CPUE

were not (Table 3, 4). The latter two were not significant because brook trout were relatively scarce in this stream but relative proportions were high given very few other fish species at the upstream sites. Brown trout density was significantly higher in the downstream sections but CPUE and relative proportions were not. This species was more prevalent in the downstream sections most likely a result of not being able to move upstream during low water conditions and from patterned stocking by state agencies into the mainstream Manistee River. Waters (1999) showed that brook trout declined after sedimentation and then continued to decline while brown trout increased. Brown trout abundances at this location were relatively low, indicating low impacts to resident brook trout populations. Also, Fausch and White (1981) showed that brook trout tend to be more abundant in headwaters and brown trout more abundant downstream as a result of competition, consumption of smaller brook trout by larger brown trout, easy catchability of brook trout by angling, and physical stream changes.

Burbot and Chinook salmon density, CPUE, and relative proportions were all significantly higher in the downstream sections. Burbot were not able to move upstream above the perched culvert. They were located generally in undercut banks in the downstream reaches. Chinook salmon juveniles were not able to move upstream past the perched culvert, at least during base flow conditions. Coho salmon and johnny darter density and CPUE were significantly higher in the downstream sections while relative proportions of each were not. These species both could not move upstream past the perched culvert but could readily move in from the mainstream Big Manistee River possibly for refuge, food, and cooler water as Sickle Creek was always cooler than the mainstream. Beschta and Taylor (1988) suggested that stream temperature in many

regions has increased as a result of land use practices, providing increased sunlight and warmer temperatures on coldwater species. Sickle Creek, however, has large quantities of surrounding riparian vegetation with full cover near the confluence. This refuge from warmer temperatures likely provides a thermal refuge for some fish.

Mottled sculpin relative proportions were significantly higher in the upstream sections but density and CPUE were not. It is unknown how long the culvert has been perched. Also the only time fish can pass the perched culvert in the upstream direction would be during high discharge intervals when water velocity would be quite high. Western blacknose dace, brook stickleback, central mudminnow, northern redbelly dace, and rainbow trout did not have any significant differences at this site. Most of these species were present in very small numbers. Longnose dace and white sucker were absent from this site, although periodic usage is possible from migrating fish from the mainstream river.

Individual Species at Milks Road, Bear Creek: Brook trout density was the only measurement at this site that was significantly higher in the upstream sections (Tables 3, 4). Brook trout predominantly reside in headwater areas of streams in the Midwest including this one (Wootton 1998). Optimal water quality, substrate, discharge, less competition with other species, and food availability help to explain this pattern. In similar fashion, Alexander and Hansen (1988) suggested that brook trout populations would increase after declining from an experimental addition of sand in Hunt Creek, Michigan. The results showed initially that the brook trout population declined to half of the pre-sand abundance. Age-0 brook trout had reached their apparent maximum

productivity ten years after sand abatement measures were implemented (namely sediment traps), indicating other possible influences. Adult brook trout have nearly completely recovered. In contrast, Curry and MacNeil (2004) showed that groundwater inputs may alleviate sedimentation in spawning areas and increase survival and durability of salmonids. Zones where groundwater inputs are large may allow quite speciose food webs to exist (Stanford and Ward 1993).

Western blacknose dace, brook stickleback, brown trout, burbot, central mudminnow, Chinook salmon, coho salmon, creek chub, johnny darter, longnose dace, mottled sculpin, northern brook lamprey, northern redbelly dace, rainbow trout, and white sucker did not have significant differences. Culverts at this location were larger and fish passage was never a problem. Human disturbance is also greatest at this site in contrast to the other sites.

NMDS: Results from the multivariate analysis suggested that in some cases the fish assemblages separated differently above vs. below a restoration site, but in others there was no apparent separation. Both the USFS 5575 (Sickle Creek) and Pine Lake Road (Pine Creek, Figure 7) sites showed separation among the fish assemblages between upstream and downstream sites, while the Steinberg Road (Pine Creek, Figure 8) and Milks Road (Bear Creek) sites did not show the same separation between fish assemblages (Table 5). At the USFS 5575 (Figure 9) site, two major gradients captured most of the variance in the fish communities, the first two dimensions containing 35.2% and 37.0%, respectively, of the information in the analytical data set. Similarly, at the Pine Lake Road site, two major gradients captured most of the variance in the fish

communities, 47.5% and 46.2%, respectively. At the Milks Road (Figure 10) site, the first two dimensions only captured 20.6% and 20.5%, respectively, while at the Steinberg Road had less than 1.0% of the variance captured by the first two dimensions. In each NMDS output, each higher dimension improved the model very little and, at most, a two-dimensional solution was recommended.

Perched culverts at both the USFS 5575 and Pine Lake Road sites were the likely explanation as to why the upstream and downstream fish assemblages separated differently on the NMDS plots. At the Steinberg Road and Milks Road sites, the culverts were not similarly perched, thus not changing the upstream to downstream fish assemblages at each site. However it is important to note that the degree to which the culverts were perched was much greater in Sickle Creek. At Pine Lake Road, dual culverts were not perched, but did experience significantly higher velocity vs. the natural stream channel.

Conclusion:

The restoration that has since occurred at these sites will likely have a dramatic effect on the fish assemblage given that we have documented some differences above and below the road stream crossings prior to restoration. Indeed, ongoing monitoring conducted by GVSU graduate students Kristofor Nault (personal communication) and Jason DeBoer (DeBoer 2008) has documented significant post-restoration effects.

Discontinuity caused by a perched culvert led to higher downstream species diversity, density and CPUE in a small tributary stream. Impacts of road-stream crossings in larger streams had multiple negative effects both up and downstream. Higher

brown and rainbow trout downstream at Pine Lake and Steinberg Road (Pine Creek) indicated that negative impacts are not always felt downstream of sediment sources. Larger substrate, larger woody debris, deeper pools, water quality, food preference, and discharge may be involved. We believe that improperly designed culverts that are too small may lead to poor conveyance of floodwaters and decrease upstream habitat quality. As was done in this project, improvements to road-stream crossings and bank stabilization should be done in such a way as to maximize and restore natural stream structure and function mainly through a return to a more natural flow regime that accommodates floods and sediment transport.

Literature Cited:

- Alexander, G.R., and E.A. Hansen. 1988. Decline and recovery of a brook trout stream following an experimental addition of sand sediment. Michigan Department of Natural Resources Fisheries Research Report No. 1943.
- Allan, J.D. 2004. landscapes and riverscapes: the influence of land use on stream ecosystems. Annual Review of Ecological Evolutionary Systematics. Vol 35: 257-84.
- Allan, J.D. and A. S. Flecker. 1993. Biodiversity conservation in running waters. Bioscience. Vol. 43(1): 32-43.
- Allen, M., Combs, D.L., Cook, S.B., and M.R. Edwards. 2003. Comparison of single-pass electrofishing to depletion sampling for surveying fish assemblages in small warmwater streams. Journal of Freshwater Ecology. Vol 18, no. 4:625-634.
- Barbour, M.T., Gerritsen, J., Snyder, B.D., and J.B. Stribling. 1999. Rapid bioassessment protocols for use in wadeable streams and rivers. United States Environmental Protection Agency. 2nd Ed.
- Barton, B.A. 1977. Short-term effects of highway construction on the limnology of a small stream in southern Ontario. Freshwater Biology: 7, 99-108.

- Berg, L., and T.G. Northcote. 1985. Changes in territorial, gill flaring, and feeding behavior in juvenile coho salmon, following short-term pulses of suspended sediment. Canadian Journal of Fisheries and Aquatic Sciences: 42, 1410-1417.
- Bernhardt, E.S., Palmer, M.A., Allan, J.D., Alexander, G., Barnas, K., Brooks, S., Carr, J., Clayton, S., Dahm, C., Follstad-Shah, J., Galat, D., Gloss, S., Goodwin, P., Hart, D., Hassett, B., Jenkinson, R., Katz, S., Kondolf, G.M., Lake, P.S., Lave, R., Meyer, J.L., O'Donnell, T.K., Pagano, L., Powell, B., and E. Sudduth. 2005. Synthesizing U.S. river restoration efforts. Science. Vol. 308: 636-637.
- Beschta, R.L. 1978. Long-term patterns of sediment production following road construction and logging in the Oregon Coast Range. Water Resources Research: 14, 1011-1016.
- Beschta, R.L., and R.L. Taylor. 1988. Stream temperature increases and land use in a forested Oregon watershed. Water Research Bulletin, 24:19-25
- Bjornn, T.C. 1971. Trout and salmon movements in two Idaho stream as related to temperature, food, streamflow, cover, and population density. Transactions of the American Fisheries Society: 100, 423-438.
- Brazner, J.C., Tanner, D.K., Detenbeck, N.E., Batterman, S.L., Stark, S.L., Jagger, L.A., and V.M. Snarski. 2005. Regional, watershed, and site-specific environmental influences on fish assemblage structure and function in western Lake Superior tributaries. Canadian Journal of Fisheries and Aquatic Sciences, 62: 1254-1270
- Carline, R.F. 2001. Effects of high-frequency pulsed-DC electrofishing on a wild brown trout population. North American Journal of Fisheries Management: 21, 571-579.
- Childers, W.F., Heckrotte, C., and R.W. Larimore. 1959. Destruction and reestablishment of stream fish and invertebrates affected by drought. Transactions of the American Fisheries Society. 88:261-285.
- Chisholm, J.L., and S.C. Downs. 1978. Stress and recovery of aquatic organisms as related to highway construction along Turtle Creek, Boone County, West Virginia. US Geological Survey Water Supply Paper 2055, Washington DC.
- Cline, L.D., Short, R.A., Ward, J.V., Carlson, C.A., and H.L. Gary. 1983. Effects of highway construction on water quality and biota in an adjacent Colorado mountain stream. U.S. Forest Service Research Note RM-429.
- Curry, R.A., and W.S. Macneill. 2004. Population-level responses to sediment during early life in brook trout. Journal of the North American Benthological Society: 23(1), 140-150.

- De Lange, H.J., De Jonge, J., Den Besten, P.J., Oosterbaan, J., and E.T.H.M. Peeters. 2004. Sediment pollution and predation affect structure and production of benthic macroinvertebrate communities in the Rhine-Meuse delta, the Netherlands. Journal of the North American Benthological Society. 23(3):557-579.
- Doonan, C.J., and G.E. Hendrickson. 1972. Reconnaissance of the Manistee River, a cold-water river in the northwestern part of Michigan's southern peninsula: U.S. Geological Survey Hydrologic Investigations Atlas 346, 2 sheets, scale 1:62,500.
- Elwood, J.W., and T.F. Waters. 1969. Effects of floods on food consumption and production rates of a stream brook trout population. Transactions of the American Fisheries Society. 98:253-262.
- Fausch, K.D., and R.J. White. 1981. Competition between brook trout and brown trout for positions in a Michigan stream. Canadian Journal of Fisheries and Aquatic Sciences: 38, 1220-1227.
- Fausch, K.D., Torgerson, C.E., Baxter, C.V., and W.L. H.W. Li. 2002. Landscape to riverscapes: bridging the gap between research and conservation of stream fishes. Bioscience. Vol 52(6): 483-498.
- Harding, J.S., Benfield, E.F., Bolstad P.V., Helfman, G.S., and E.B.D. Jones III. 1998. Stream biodiversity: the ghost of land use past. Proceedings from the National Academy of Sciences. Vol. 95: 14843-14847.
- Helms, B.S., Schoonover, J.E., and J.W. Feminella. 2008. Assessing influences of hydrology, physiocochemistry, and habitat on stream fish assemblages across a changing landscape. Journal of the American Water Resources Association. Vol 45(1): 157-69.
- Jacobson, R.B. and D.L. Galat. 2008. Design of a naturalized flow regime an example from the Lower Missouri River, USA. Ecohydrology. Vol 1: 81-104.
- Jansson, R., Backx, H., Boulton, A.J., Dixon, M., Dudgeon, D., Hughes, F.M.R., Nakamura, K., Stanley, E.H., and K. Tockner. 2005. Stating mechanisms and refining criteria for ecologically successful river restoration: a comment on Palmer et al. (2005). Journal of Applied Ecology. Vol 42: 218-222.
- Jones, E.B.D. III, Helfman, G.S., Harper, J.O., and P.V. Bolstad. 1999. Effects of riparian forest removal on fish assemblages in southern Appalachian streams. Conservation Biology. Vol 13, no. 6: 1454-1465.
- Jungwirth, M., Muhar, S., and S. Schmutz. 2002. Re-establishing and assessing ecological integrity in riverine landscapes. Freshwater Biology. Vol 47: 867-887.

- Junk, W.J., Bayley, P.B., and R.E. Sparks. 1989. The flood pulse concept in river-floodplain systems. Proceeding of the International Large River System Symposium, Canadian Fisheries and Aquatic Sciences.
- Kruskal, J.B. 1964. Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. Psychometrika 29:1-27.
- Lake, P.S. 2000. Disturbance, patchiness, and diversity in streams. Journal of the North American Benthological Society: 19(4), 573-592.
- Larimore, R.W., Childers, W.F., and C. Heckrotte. 1959. Destruction and reestablishment of stream fish and invertebrates affected by drought. Transactions of the American Fisheries Society: 88, 261-285.
- Lisle, Thomas E. 1989. Sediment transport and resulting deposition in spawning gravels, north coastal California. Water Resources Research: 25(6), 1303-1319.
- Lisle, T., and S, Hilton. 1991. Fine sediment in pools: an index of how sediment is affecting a stream channel. United States Forest Service R-5 Fish Habitat Relationship Technical Bulletin No. 6.
- Loomis, J.B. 1989. A bioeconomic approach to estimating the economic effects of watershed disturbance on recreational and commercial fisheries. Soil and Water Conservation: 44(1), 83-87.
- Mann, R.H.K., and T. Penczak. 1986. Fish production in rivers: a review. Polskie Archiwum Hydrobiologii, 33:233-247
- Mather, P.M. 1976. Computational methods of multivariate analysis in physical geography. J. Wiley & Sons, London. 532 pp.
- Maunder, M.N., Sibert, J.R., Fonteneau, A., Hampton, J., Kleiber, P., and S.J. Harley. 2006. Interpreting catch per unit effort data to assess the status of individual stocks and communities. ICES Journal of Marine Science. 63:1373-1385.
- McCune, B., and J.B. Grace. 2002. Analysis of ecological communities. MjM Software Design, Gleneden Beach, Oregon.
- Meyer, J.L., Strayer, D.L., Wallace, J.B., Eggert, S.L., Helfman, G.S., and N.E. Leonard. 2007. The contribution of headwater streams to biodiversity in river networks. Journal of the American Water Resources Association. Vol 43(1): 86-103.
- Natural Resources Conservation Service. 2004. Landforms of Northern Lower Michigan: NRCS Michigan State OMCE.

- Palmer, M.A., Bernhardt, E.S., Allan, J.D., Lake, P.S., Alexander, G., Brooks, S., Carr, J., Clayton, S., Dahm, C.N., Follstad Shah, J., Galat, D.L., Loss, S.G., Goodwin, P., Hart, D.D., Hassett, B., Jenkinson, R., Kondolf, G.M. Lave, R., Meyer, J.L., O'Donnell, T.K., Pagano, L., and E. Sudduth. 2005. Standards for ecologically successful river restoration. Journal of Applied Ecology. Vol 42: 208-217.
- Peters, J.L. 1967. Effects on a trout stream of sediment from agricultural practices. Journal of Wildlife Management. 31:805-812.
- Saunders, J.W., and M.W. Smith. 1965. Changes in a stream population of trout associated with increased silt. Journal of the Fisheries Research Board of Canada. 22:395-404.
- Schlosser, I.J. 1991. Stream fish ecology: a landscape perspective. Bioscience. Vol 41(10): 704-711.
- Scott, J.B., Steward, C.R., and Q.J. Stober. 1986. Effects of urban development on fish population dynamics in Kelsey Creek, Washington. Transactions of the American Fisheries Society: 115, 555-567.
- Seelbach, P.W. and M.J. Wiley. 1997. Overview of the Michigan Rivers Inventory (MRI) Project. Michigan Department of Natural Resources Fisheries Division Technical Report Number 97-3.
- Servizi, D.A., and D.W. Martens. 1992. Sublethal responses of coho salmon to suspended sediments. Canadian Journal of Fisheries and Aquatic Sciences: 49, 1389-1395.
- Stanford, J.A., and J.V. Ward. 1993. An ecosystem perspective of alluvial rivers: connectivity and the hyporheic corridor. Journal of the North American Benthological Society, 12(1): 48-60
- Torgersen, C.E., Baxter, C.V., Li, H.W., and B. A. McIntosh. 2006. Landscape influences on longitudinal patterns of river fishes: spatially continuous analysis of fish-habitat relationships. American Fisheries Society, 48:473-492.
- United States Geological Survey. 1990. Land Cover Manistee County: Michigan State University NCCD, Michigan State University Board of Trustees.
- Ward, J.V., Tockner, K., Uehlinger, U., and F. Malard. 2001. Understanding natural patterns and processes in river corridors as the basis for effective river restoration. Regulated Rivers: Research & Management. Vol 17: 311-323.
- Wang, L., Lyons, J., Rasmussen, P., Seelbach, P., Simon, T., Wiley, M., Kanehi, P., Baker, E., Niemala, S., Stewart, P.M. 2003. Watershed, reach, and riparian influences on stream fish assemblages in the Northern Lakes and Forest

- Ecoregion, USA. Canadian Journal of Fisheries and Aquatic Sciences. Volume 60: 491-505.
- Waters, T.F. 1992. Annual production, production/biomass ratio, and the ecotrophic coefficient for management of trout in streams. North American Journal of Fisheries Management, 12:34-39
- Zorn, T.G., Seelbach, P.W., and M.J. Wiley. 1998. Patterns in the distributions of stream fishes in Michigan's lower peninsula. Michigan Department of Natural Resources Fisheries Division Research Report Number 2035.

Tables:

Table 1. Mean water quality variables (+1 SD) for each respective reach sampled. Each reach was sampled twice each year, May/June and late July/August 2004 and 2005, except for Bear Creek, which had one sample period in 2004, and two in 2005. N.A. = not available.

River	Abbrev.	7. GPS Coordinates		Mean		Mean	Mean	Mean Specific	Mean
		Start	End	Temperature (0°C)	ature	pН	Dissolve Oxygen (mg/L)	Conductivity (mS/cm ²)	Discharge (m³/s)
Pine	HUFF	N 44º 14' 43.4"		14.6 (0.	53)	8.42	14.8	0.28 (0.01)	52.3 (23.0)
		W086º 03' 41.2"	Bridge	`	,	(0.02)	(4.19)	` ′	` ′
	STDS3	N 44º 13' 45.7"	N 44º 13' 41.9"	13.8 (0.	40)	8.05	10.1	0.41 (0.15)	28.0 (4.68)
		W086° 01' 49.9"	W086° 01' 43.7"			(0.29)	(1.31)		
	STDS2	N 44º 13' 40.2"	N 44º 13' 36.6"	13.8 (0.	40)	8.05	10.1	0.41 (0.15)	28.0 (4.68)
		W086° 01' 40.3"	W086° 01' 40.3"			(0.29)	(1.31)		
	STDS1	N 44º 13' 34.9"		13.8 (0.	40)	8.05	10.1	0.41 (0.15)	28.0 (4.68)
		W086° 01' 36.1"	Bridge			(0.29)	(1.31)		
	STUS1		N 44º 13' 26.3"	13.9 (1.	26)	8.14	10.4	0.42 (0.16)	26.3 (15.2)
		Bridge	W086° 01' 23.0"			(0.16)	(0.47)		
	STUS2	N 44º 13' 24.4"	N 44º 13' 19.3"	13.9 (1.	26)	8.14	10.4	0.42 (0.16)	26.3 (15.2)
	******	W086° 01' 20.3"	W086º 01' 23.5"			(0.16)	(0.47)	0.42.40.4.5	
	USFS 8430	N.A.	N.A.	14.1 (0.	84)	8.23	10.4	0.42 (0.16)	26.1 (15.1)
	DI D.GO		N. 440 401 00 48 N400 50	1450		(0.22)	(0.50)	0.42 (0.14)	22.0 (10.2)
	PLDS3	N 44º 12' 34.5"	N 44° 12′ 36.1″ W085°	14.5 (0.	/1)	8.13	11.2	0.42 (0.14)	23.0 (10.2)
	PLDS2	W085° 59' 11.1"	59' 03.5" N 44º 12' 38.5" W085º	145 (0)	71)	(0.22)	(0.85)	0.42 (0.14)	22.0 (10.2)
	PLDS2	N 44º 12' 37.9" W085º 58' 00.8"	58' 53.3"	14.5 (0.	/1)	8.13	11.2 (0.85)	0.42 (0.14)	23.0 (10.2)
	PLDS1	N 44º 12' 37.6"	56 53.3	14.5 (0.	71)	(0.22) 8.13	(0.85)	0.42 (0.14)	22.0 (10.2)
	PLDSI	W085° 58' 49.0"	Bridge	14.5 (0.	/1)	(0.22)	(0.85)	0.42 (0.14)	23.0 (10.2)
	PLUS1	VV003 30 49.0	N 44º 12' 34.0"	14.4 (0.	73)	8.17	11.3	0.42 (0.14)	20.2 (11.0)
	I LUSI	Bridge	W085° 58' 35.9"	14.4 (0.	13)	(0.19)	(0.63)	0.42 (0.14)	20.2 (11.0)
PLUS2	N 44º 12' 34.9"	N 44º 12' 33.8"	14.4 (0.	73)	8.17	11.3	0.42 (0.14)	20.2 (11.0)	
	12002	W085° 58' 32.2"	W085º 58' 24.8"	1 (0.	,,,	(0.19)	(0.63)	0.12 (0.11)	20.2 (11.0)
Sickle	SIDS3	N 44º 17. 632'	N 44° 17. 678' W086°	13.9 (0.	51)	8.57	11.1	0.41 (0.16)	2.88 (0.72)
		W086º 08.929'	08.957'		- /	(0.09)	(0.39)	(3.3)	,
	SIDS2	N 44° 17' 41.7"	N 44º 12' 38.5" W085º	13.9 (0.	51)	8.57	11.1	0.41 (0.16)	2.88 (0.72)
		W086º 09' 06.6"	58' 53.3"			(0.09)	(0.39)		
	SIDS1	N 44º 17' 42.0"		13.9 (0.	51)	8.57	11.1	0.41 (0.16)	2.88 (0.72)
		W086º 09' 13.5"	Bridge			(0.09)	(0.39)		
	SIUS1		N 44º 17' 47.6"	13.7 (0.	13)	8.60	11.2	0.42 (0.14)	2.07 (0.94)
		Bridge	W086° 09' 14.2"			(0.12)	(0.53)		
	SIUS2	N 44º 17' 49.8"	N 44º 17' 50.4"	13.7 (0.	13)	8.60	11.2	0.42 (0.14)	2.07 (0.94)
		W086° 09' 16.9"	W086º 09' 20.0"			(0.12)	(0.39)		
Bear	LOWER	N 44º 17' 22.4"	N 44° 17' 27.6"	16.2 (1.	05)	8.56	11.3	0.29 (0.04)	144 (80.6)
	BEAR	W086° 03' 46.6"	W086º 04' 01.9"			(0.34)	(2.55)		
	SPIRIT	N 44º 18' 42.0"		15.4	8.45	14.2	(6.58)	0.29 (0.04)	279 (203)
	OF	W086° 02'	Deider	(0.59)	(0.13)				
	WOODS	56.9"	Bridge	165	0.52	12.0	(6.00)	0.28 (0.02)	226 (146)
	JOHNSON	N.A.	N.A.	16.5	8.53	15.9	(6.90)	0.28 (0.02)	236 (146)
	SWAINDS2	N 44º 20' 56.6"	N 44º 21' 03.0"	(1.09) 18.5	(0.17) 8.68	15.6	(7.11)	0.31 (0.03)	271 (125)
	5 WAIND 52	W086º 03' 09.9"	W086º 03' 06.7"	(2.16)	(0.30)	13.0	(7.11)	0.51 (0.05)	2/1 (123)
	SWAINDS1	N 44º 21' 02.4"	N 44º 21' 09.6"	18.5	8.68	15.6	(7.11)	0.31 (0.03)	271 (125)
	SWAINDSI	W086º 03' 09.8"	W086º 03' 06.0"	(2.16)	(0.30)	13.0	(7.11)	0.51 (0.05)	2/1 (123)
	MILKSDS	N.A.	N.A.	17.7	8.64	113	(1.44)	0.33 (0.02)	217 (63.0)
	111111111111111111111111111111111111111	. 1./ 1.	. 1 2.	(1.60)	(0.25)	11.3	(1.17)	0.05 (0.02)	217 (03.0)
	MILKSUS	N.A.	N.A.	17.7	8.64	11.3	(1.44)	0.33 (0.02)	217 (63.0)
			··· 	(1.60)	(0.25)		,	· · · · · · · · · · · · · · · · · · ·	(0)

Table 2. Average abundance data for most influential species affecting site differences in NMDS. The three most influential species were selected and their average abundance per reach in each respective reach has been given.

Reach and Average Number/Reach						
Site	Species	US2	US1	DS1	DS2	DS3
Pine Lake Road (Pine	Brook stickleback	4	7	0	0	0
Creek)	Rainbow trout	0	1	5	7	5
	White sucker	5	5	3	0	0

Steinberg Road	Brown trout	3	2	5	5	13
(Pine Creek)	Mottled sculpin	8	12	7	12	41
	Rainbow trout	3	4	5	8	11
USFS 5575	Burbot	0	0	15	12	8
(Sickle Creek)	Chinook	0	0	3	3	2
	salmon					
	Coho salmon	0	0	4	4	6
		US	DS	SWAINUS	SWAINDS	
Milks Road	Brown trout	11	8	7	6	
(Bear Creek)	Mottled sculpin	29	19	13	9	
	Rainbow trout	71	52	47	44	

Table 3. Mean density, CPUE, and relative abundance data for significant differences between upstream and downstream locations at respective sites. Standard deviations are in parentheses. Abbreviations are as follows: WBND = western blacknose dace, BRS = brook stickleback, BRT = brook trout, BNT = brown trout, BUT = burbot, CEM = central mudminnow, CHS = Chinook salmon, COH = coho salmon, CRC = creek chub, JOD = Johnny darter, MOS = mottled sculpin, RBT = rainbow trout, WHS = white sucker.

	Mean Relative Density (ha)			Mean CPUE		Mean Relative Abundance		ce	
Species and Location	Upstream	Downstream	p-value	Upstream	Downstream	p-value	Upstream	Downstream	p-value
WBND (Pine Lake Rd.)	430 (60)	920 (200)	0.002	2.8 (0.64)	5.11 (0.73)	0.002	24.8 (2.0)	46.6 (3.18)	0.001
WBND (Steinberg Rd.)							60.0 (4.0)	48.5 (4.1)	0.05
BRS (Pine Lake Rd.)	55 (2.0)	3.0 (1.0)	0.001	0.38 (0.15)	0.013 (0.008)	0.001	3.17 (1.1)	0.09 (0.05)	0.001
BRT (Milks Rd.)	6.0 (4.0)	1.0 (0.05)	0.031						
BRT (USFS 5575)							3.19 (1.5)	0.18 (0.18)	0.035
BNT (USFS 5575)	5.0 (7.0)	37 (2.0)	0.046						
BNT (Steinberg Rd.)	30 (7.0)	80 (7.0)	0.049				3.56 (0.76)	7.45 (1.6)	0.05
BUT (USFS 5575)	0	45 (20)	0.01	0	0.84 (0.41)	0.01	0	13.6 (3.4)	0.002
CEM (Pine Lake Rd.)	260 (60)	62 (3.0)	0.001	1.58 (0.38)	0.37 (0.17)	0.001	13.8 (1.6)	3.06 (0.78)	0.001
CHS (USFS 5575)	0	103 (50)	0.013	0	0.19 (0.09)	0.001	0	4.15 (1.1)	0.004
COH (USFS 5575)	0	18.3 (10)	0.04	0	0.31 (0.06)	0.036			
CRC (Pine Lake Rd.)							16.9 (1.5)	12.2 (1.5)	0.033
JOD (USFS 5575)	0	70 (4.0)	0.02	0	0.153 (0.10)	0.049			
MOS (Pine Lake Rd.)	13 (1.0)	31 (10)	0.008	0.009 (0.01)	0.20 (0.10)	0.017	0.066 (0.06)	1.89 (0.68)	0.023
MOS (USFS 5575)							85.6 (4.0)	54.8 (5.7)	0.001
RBT (Pine Lake Rd.)	50 (4.0)	5.4 (2.0)	0.001	0.04 (0.03)	0.34 (0.12)	0.002	0.294 (0.16)	3.43 (1.1)	0.016
RBT (Steinberg Rd.)	38 (1.0)	80 (2.0)	0.031				4.91 (0.74)	7.66 (0.89)	0.028
WHS (Pine Lake Rd.)	50 (20)	10 (6.0)	0.001	0.31 (0.11)	0.060 (0.04)	0.002	2.65 (0.60)	0.429 (0.29)	0.001

Table 4. Summary of various taxa-specific and community level responses above and below restoration sites.

	Undersized culverts			Bridge/bank erosion
Observed response	Pine Lake Road (Pine	Steinberg Road (Pine	USFS 5575 (Pine	Milks Road (Bear
	Creek)	Creek)	Creek)	Creek)
Higher upstream	Brook stickleback	Western blacknose	Brook trout	Brook trout
	Central mudminnow	dace	Mottled sculpin	
	White sucker			
Higher downstream	Western blacknose	Rainbow trout	Brown trout	
	dace			
	Mottled sculpin	Brook trout	Burbot	
	Rainbow trout		Chinook salmon	
			Coho salmon	
			Johnny darter	
Absent	Burbot	Longnose dace		
	Chinook salmon	Northern brook		
	Northern brook	lamprey		
	lamprev			

Table 5. Set-up and results of NMDS analysis for each restoration site.

	Pine Lake Road (Pine Creek)	Steinberg Road (Pine Creek)	USFS 5575 (Sickle Creek)	Milks Road (Bear Creek)
Distance measure	Sorenson	Sorenson	Sorenson	Sorenson
Software used	PCORD	PCORD	PCORD	PCORD
Starting coordinates	Random	Random	Random	Random
Reduction in dimensionality	1	1	1	1
Step length	0.2	0.2	0.2	0.2
Random number of seeds	Use time	Use time	Use time	Use time
Run with real data	40	40	40	40
Runs with randomized data	50	50	50	50
Autopilot	Yes	Yes	Yes	Yes
Dimensionality	Scree plot	Scree plot	Scree plot	Scree plot
Number of axes	1	Not found	1	Not found
Maximum iterations	400	400	400	400
Final stress	0.15276	Not found	0.00001	Not found
Final iterations for best solution	49	Not found	31	Not found
Stability criterion	0.00001	0.00001	0.00001	0.00001
R-squared	0.937	0.001	0.722	0.416
Monte Carlo Test (p)				
Number of axes]			
1	0.0196	0.549	0.0392	0.3922
2	0.3922	0.7451	0.4118	0.9804
3	0.8235	0.9412	0.9412	1
4	0.9608	0.9608	1	1
5	0.9608	0.9608	1	1
6	0.9804	0.9608	1	1

Figures:




Figure 1. Sickle, Pine, and Bear Creeks within the Manistee River Watershed.




Figure 2. Sickle Creek site design.




Figure 3. Total fish density above and below road stream crossings. Pine Lake and Steinberg Road = Pine Creek, USFS 5575 = Sickle Creek, Milks Road = Bear Creek. Standard error bars are also present. Sites are listed on the x-axis. Reach, season and year are pooled; Pine Lake = 8 & 12 (up/dn), Steinberg = 8 & 12 (up/dn), USFS = 8 & 12 (up/dn), & Milks = 3 & 9 (up/dn).




Figure 4. Fish species dominance (Simpson's Index) above and below road stream crossings. Standard error bars are also present. Sites are listed on the x-axis. Data pooled and sample size as in Fig. 3.




Figure 5. Species diversity (Shannon's Diversity) above and below road stream crossings. Standard error bars are also present. Sites are listed on the x-axis. Data pooled and sample size as in Fig. 3.




Figure 6. Species richness above and below road stream crossings. Standard error bars are also present. Sites are listed on the x-axis. Data pooled and sample size as in Fig. 3.

Pine Lake Road Crossing




Figure 7. Fish assemblage differences among Pine Lake Road reaches, Pine Creek, based on NMDS analysis. Site abbreviations as in Table 1 and set-up and results in Table 5.

Steinberg Road Crossing

Fish composition (relative abundance (m2)




Figure 8. Fish assemblage differences among Steinberg Road reaches, Pine Creek. Site abbreviations as in Table 1 and set-up and results in Table 5.

USFS 5575 Road Crossing

Fish composition (relative abundance (m²))




Figure 9. Fish assemblage differences at Sickle Creek (USFS road 5575). Site abbreviations as in Table 1 and set-up and results in Table 5.

Milks Road Crossing

Fish composition (relative abundance (m²))




Figure 10. Fish assemblage differences among Milks Road reaches, Bear Creek. Site abbreviations as in Table 1 and set-up and results in Table 5.

CHAPTER 3

LONGITUDINAL FISH DISTRIBUTION IN THREE WEST MICHIGAN STREAMS OF DIFFERENT ORDERS

ABSTRACT

Few studies have examined fish longitudinal distribution in river systems impaired by high rates of sedimentation, or, for that matter, in the Great Lakes Region. The purpose of this study was to 1) provide a detailed description of the fish assemblage structure from headwaters to mouth on the three study streams, and 2) assess the differences between streams regarding fish assemblage structure. The study included three tributary streams (Sickle Creek, 1st order, Pine Creek, 2nd order, Bear Creek, 4th order) within the lower Manistee River watershed. A total of 24 electrofishing reaches comprised the longitudinal gradient analysis. Longitudinal analysis and non-metric multidimensional scaling showed unique assemblages between Sickle, Pine, and Bear Creeks. Sickle Creek was most uniquely different with occurrence of brook trout in headwater reaches and burbot in downstream reaches; a pattern attributed to the presence of a largely impassable perched culvert that separated up from downstream reaches. Pine Creek had northern redbelly dace throughout, whereas neither of the other two streams had this species present. This is likely due to the colder thermal regime of Pine Creek, an environmental attribute preferred by northern redbelly dace. Pine Creek was also characterized by an abundance of brown and rainbow trout in the most downstream location (Huff Road) again due to cold thermal regime and abundance of large woody debris. Bear Creek was characterized by burbot only in downstream sections, as well as longnose and western blacknose dace, and an increase in brook trout upstream. As such, Bear Creek conformed to the more expected patterns in longitudinal distribution. In conclusion, different order streams and patterns seemed to reflect the environmental habitat template, even when this template deviated from expected conditions.

Introduction:

Fish longitudinal distribution has been shown to be affected by biotic and abiotic factors. For example, Grenouillet et al. (2004) studied within basin distribution of local species richness in the Upper Saone River, France, and found that stream width and gradient significantly influenced local species richness (LSR). Schaefer and Kerfoot (2004) working in Illinois showed a correlation between distance from the mouth of the

stream and assemblage variability over time, as well as a negative correlation between distance from the mouth of the stream and mean diversity over time. They hypothesized that the observed patterns in community variability and the distribution of some species within the drainage are best explained by the interactions between the fauna of big and small rivers. In their study, Grenouillet et al. (2004) showed that spatial factors also influenced the within-basin distribution of LSR and resulted in spatial autocorrelation, highlighting biotic processes in structuring stream fish assemblages. However, the study did not confirm other published predictions that headwater streams entering large rivers directly should have greater species richness. The spatial autocorrelation was only significant in larger rivers (from 4th- to 7th-order streams), suggesting that the relative importance of local habitat and biotic processes may depend on the position along the longitudinal gradient.

Some studies have examined correlations between stream order and fish assemblage measurements within the United States. Paller (1994) examined relationships between fish assemblage structure and stream order in South Carolina coastal plain streams. Average species richness adjusted to a constant stream surface area were 12.7, 17.5, 21.4, and 22.0 species in first- through fourth-order streams, respectively. Species addition and replacement led to large changes in species composition among stream orders. Relatively small fishes numerically dominated headwater streams. Relatively large fishes were most common in fourth-order streams. Headwater species richness was lower and longitudinal species replacement was greater than often observed in other geographic regions of the United States. A comparative assessment of long-term temperature and precipitation records suggested that high species richness at headwater

sites was related to mild climate and lack of steep elevation gradients. The presence of numerous small headwater species created the potential for multiple species replacements as downstream increases in habitat volume permitted the establishment of larger fish with predatory and competitive advantages. Smith and Kraft (2005) showed that the proportion of fine substrate, canopy cover, in-stream vegetation, and water temperature were the four local habitat factors related to the abundance of fish species in the Beaverkill- Willowemoc watershed in New York. Confluence link and stream order were the stream network position measures with the greatest influence on fish assemblages. The results showed that stream fish assemblages in the study watershed were influenced by a combination of small-scale habitat variables and stream position within a watershed network.

Only a handful of studies have examined fish longitudinal distribution in river systems impaired by high rates of sedimentation, or, for that matter, in the Great Lakes Region (Seelbach and Wiley 1997, Zorn et al. 1998, Fausch et al. 2002). For example, Thomas (2002) examined fish and invertebrate communities and habitat of the fifth-order mainstem, two second-order adventitious tributaries to the mainstem, and three second-order headwater streams of the Pine River (Alcona County, Michigan) from May to August 2000. Fish species richness generally increased with increasing stream order and was higher in the adventitious streams than in the headwater streams. Little published research is evident in the Great Lakes regarding entire fish assemblages, their structure, their differences between stream order, and also the influence of barriers to fish migration. The purpose of this study was to: (1) provide a detailed description of the fish assemblage structure from headwaters to mouth on the two larger study streams, (2) to

assess fish assemblage structure above and below a perched culvert, and (3) assess the differences in fish assemblage structure between streams.

Materials and Methods:

Fish Longitudinal Distribution: Fish assemblage relative abundance data were used to construct stacked area plots with respective sites in order from upstream to downstream in each season. Basic descriptive data trends were inferred using patterns visible from stacked area plots. Species catch/minute data were compared along the gradient using ANOVA (α < 0.5).

Fish Assemblage and Longitudinal Analyses: Fish assemblage data were collected during spring and fall of 2004 and 2005 to determine fish longitudinal distribution between seasons and streams. Bear Creek data for spring 2004 was not collected because of high water and equipment problems. Fish assemblages were monitored along longitudinal gradients for all streams using temperature, pH, dissolved oxygen, specific conductivity, and discharge as indicators for the longitudinal fish distribution patterns. The statistical program, PCORD, and NMDS were used to evaluate the effects of these abiotic factors on fish distribution patterns, most importantly to depict which factors were affecting the fish assemblages the most with respect to the longitudinal distribution of sites in each respective stream (McCune and Grace 2002).

The first data matrix in NMDS was the fish assemblage data as relative abundance and the second data matrix included the water quality parameters (pH, dissolved oxygen (mg/L), specific conductivity (umhos/cm), temperature (0 C), and discharge (ft³/s). Multiple regression analysis was used to determine significance of water

quality variables to fish assemblages. Only whole fish community assemblage measurements were used: density, richness, dominance, and diversity.

Results and Discussion:

Longitudinal predictions:

Because of the predictable dynamic and shifting nature of lotic ecosystems from headwaters to mouth (Vannote et al. 1980), we can make some predictions of the expected fish community composition along this continuum. Many studies have documented the longitudinal shift in salmonids from headwaters to mouth and have explained these patterns based largely on temperature (Smith and Kraft 2005), wetted stream area (Grenouillet et al. 2004; Paller 2004), proximity to a larger receiving river (Grenouillet et al. 2004; Schaefer and Kerfoot 2004; Thomas 2002), and measures of discharge (Grenouillet et al. 2004). Based on this physical and chemical template, we predicted that the upper reaches of both Pine and Bear Creek would consist of more brook trout, while lower reaches would see a decline in brook trout and a relative increase in brown and rainbow trout. Similarly, we expected that the entire fish community diversity would be maximized in the mid-reaches of both Pine and Bear Creeks due to the overlap of taxa from up and down-stream fish communities, respectively.

<u>Sickle Creek</u>: Longitudinal sampling and subsequent analysis of Sickle Creek was very limited and constrained to essentially one reach. Ideally, a further reach upstream from the furthest upstream reach at this site would allow comparison along the continuum from

headwaters to mouth. However, the experimental design does allow for comparisons above and below an undersized, perched culvert.

Brook trout were only seen in upstream reaches (Figure 11). Reasons for this might include competition from other species, colder water temperatures, and the presence of a perched culvert that reduced brown trout introductions. Still only about 5% of the fish assemblage at this location consisted of brook trout. Brook trout are normally located within clear, cool, well-oxygenated creeks and small to medium rivers (Page and Burr 1991). Species found only in the downstream reaches included blacknose dace, bluntnose minnows (Pimephales notatus), burbot, and brown trout. Western blacknose dace are normally located within rocky runs and pools of headwater streams (Page and Burr 1991) while bluntnose minnows are normally located within clear, rocky streams. Many of these species may migrate in and out of this stream on a short time scale. Species that were fairly uniform throughout this stream were mottled sculpin, rainbow trout, and Chinook salmon. Mottled sculpin are normally found within the headwater reaches of streams (Page and Burr 1991). These species appeared to have good food resources and habitat. Mottled sculpin were most abundant throughout the stream. The patterns seen were very similar between seasons and years. However, Chinook and coho salmon were both located in the upstream reaches in 2004 but constrained to the downstream reaches in 2005. In 2004, increased discharge rates may have allowed for increased fish passage upstream. Species contributing most to differences above vs. below the perched culverts included brook trout, burbot, northern redbelly dace, western blacknose dace, longnose dace, and rainbow trout (Table 9).

Catch Per Unit Effort: No significant differences with fish catch per minute were seen in Sickle Creek longitudinally in any reaches measured (Table 6, Figure 14). Mottled sculpin were the most dominant species in all reaches sampled and did not change significantly from headwaters to mouth.

Pine Creek: Western blacknose dace steadily increased in relative abundance from upstream to downstream reaches and then decreased steadily from U.S Forest Service Road 8430 to Huff Road, thus peak densities occurred in the mid reaches of the stream (Figure 12). This species prefers upstream sections within streams (Wootton 1998). Other reasons for this distribution may include habitat, competition, and greater abundance of predatory fish (trout) in downstream reaches. Brook trout were low in relative abundance at all reaches, a fact that may be correlated with increased sedimentation and competition with brown trout. Northern redbelly dace were more abundant in upstream reaches versus lower reaches. These species prefer colder water and usually inhabit headwater reaches of streams (Page and Burr 1991). Brown trout were more abundant at lower reaches with a large portion of the fish community at Huff Road consisting of this species. Reasons include recent stocking, proximity to the mainstream Manistee River. Furthermore, brown trout can withstand a wider range of temperatures and can inhabit a wide range of habitats (Page and Burr 1991). Rainbow trout mimicked brown trout in this respect and also increased from lower abundance in headwater reaches to a quarter of the fish community at Huff Road. These patterns were fairly consistent between seasons and are consistent with other studies examining longitudinal distribution of trout and salmonid taxa (Page and Burr 1991; Thomas 2002; Smith and Kraft 2005). Species contributing

most to differences along the river continuum included western blacknose dace, brown trout, northern redbelly dace, and rainbow trout (Table 9). Based on the higher relative abundance and CPUE of rainbow and brown trout at Huff Road we expected that the macroinvertebrate forage base would be higher at this site as well. However, De Mol (2007) found the inverse. Total macroinvertebrate abundance (m²) was significantly lower at Huff Road vs. the two upstream sites at Steinberg and Forest Service road 8430, averaging approximately 220. Unpublished gut content data suggests that terrestrial insects and smaller forage fish contribute significantly to the energy budget of the larger rainbow and brown trout (Kevin Donner and Eric Snyder, personal communication).

Catch Per Unit Effort: Western blacknose dace showed significantly higher CPUE in upstream reaches versus downstream reaches, especially the Huff Road reach where they were absent (Table 7, Figure 15). Brown trout CPUE was very high at the Huff Road reach and thus predation may explain why western blacknose dace were absent at this site. Creek chub and central mudminnow were significantly higher in upstream reaches versus downstream reaches. Northern redbelly dace were significantly more common in the upstream reaches versus downstream reaches, especially Huff Road, where they were absent. Rainbow trout were significantly higher at the Huff Road reach versus all other reaches. High rainbow and brown trout at the Huff Road reach may have negatively affected creek chub, northern redbelly dace, and central mudminnow CPUE in the same manner. Future studies should be implemented to try and explain the increased abundance of brook and rainbow trout with an apparent lack of a forage base.

Our predictions of the longitudinal distribution of the fish community in Pine Creek were generally met: higher rainbow and brown trout populations in lower Pine Creek (Huff Road) vs. upstream sites, higher abundance of northern red belly dace and western blacknose dace upstream. Interestingly, we did not observe an increase in brook trout in the upstream reaches—a fact that may simply be related to the size of the stream (2nd order), although this deserves future consideration and study.

Bear Creek: Brook trout were most abundant in upstream sections (Figure 13). Reasons could include colder water temperatures, and less competition from other species. Brook trout tend to inhabit headwater reaches of streams (Page and Burr 1991). Burbot were minimal at upstream sections and increased in lower Bear Creek, where they appeared to be located near undercut banks and sandy substrate. Most burbot were small in size, likely the result of recent spawning activity by adults. Adults spawn in clear, cool streams, while the adults reside in deep lakes (Page and Burr 1991). Most rainbow trout were present in upstream reaches versus downstream reaches, likely the result of colder stream temperatures. Western blacknose dace were more abundant in upstream sections, increasing steadily from mouth to headwaters. These species also require cooler water temperatures, but were also likely affected by predation. Species contributing most to differences along the river continuum include western blacknose dace, burbot, and longnose dace (Table 9). Fish assemblage composition was very similar between seasons.

Catch Per Unit Effort: Western blacknose dace were significantly higher in the Swain downstream reach versus the Lower Bear reach (Table 8, Figure 16). Western blacknose

dace require larger substrate and cooler water temperatures, both of which are prevalent at the Swain reach versus the downstream sites. Burbot were significantly higher in downstream reaches, mainly the lower Bear Creek reach, compared to upper reaches and may be a result of more abundant undercut banks that would be preferred by this species and proximity to the Big Manistee River. All other species were not determined to be significantly different between reaches.

Our predictions of the longitudinal distribution of the fish community in Bear Creek were similarly met: higher rainbow and brook trout populations in upper reaches vs. downstream reaches, higher abundance of western blacknose dace increasing from Lower Bear upstream, as well as higher burbot populations at Lower Bear. Interestingly, even higher populations of brook trout were found further upstream at Leffew Road (Snyder et al. 2007), thus supporting our predictions of brook trout occupying headwater reaches of streams.

Fish assemblages and water chemistry:

<u>Sickle Creek</u>: Discharge was positively correlated to and significantly explained 22.3 and 25.6% of the variation in diversity and richness, respectively, in Sickle Creek (multiple regression analyses, Figures 17 & 20). Thirty-one percent of the variation in total fish density was explained by pH. Species dominance was not significantly correlated to any water quality variable. Discharge and pH were the only significant water quality variables related to total fish density, species diversity, and species richness in Sickle Creek and showed a positive correlation. Discharge can be a good predictor of what species are in a reach, selecting for and against good versus poor swimmers, respectively (Wootton

1998), and also affecting stream wetted channel area and therefore available habitat. Total fish density in Sickle Creek was explained to some extent by pH, which is generally not thought to be a good predictor of species presence/absence, at least at the circumneutral pH values recorded in this study. This correlation is likely ecologically meaningless. Species dominance was not found to be predicted by any water quality variable.

Pine Creek: In Pine Creek, discharge explained 11% of the variation in species diversity and 15.7% of richness and may be affected similarly as mentioned for Sickle Creek (Figures 18 & 21). Species dominance and total fish density were not explained by any water quality variable measured and are likely the result of variability between sampling reaches, seasons, and interaction with biotic factors versus abiotic.

Bear Creek: In Bear Creek, temperature accounted for 23.5% of the variation in species diversity and 10.9% of dominance (Figures 19 & 22). Temperature is usually a good predictor as many species have thresholds and optimal temperatures. Brazner et al. (2005) showed that although a variety of regional, fragmentation, and storage-related factors had significant influences on the fish assemblages; water temperature appeared to be the single most important environmental factor in western Lake Superior 2nd and 3rd order tributaries. Total fish density was most strongly correlated to dissolved oxygen (24.6%) with higher dissolved oxygen levels indicative of better water quality in middle and upstream reaches, allowing for the presence of a wider range of fish species within a reach.

Bear Creek species richness was not explained by any of the water quality variables measured. Bear Creek had the highest species richness of the three sample streams and the distribution was not necessarily uniform from headwaters to mouth. Some reaches, specifically the Johnson Road reach, had the highest species richness, despite the middle position of this reach in the longitudinal continuum. Bear Creek was the largest of the streams measured had the highest species richness, and was considered to have the most stable discharge. Similarly, Taylor et al. (2006) showed the most stable assemblages were the most species-rich and occurred in relatively large, stable environments.

Conclusion:

Longitudinal analysis showed unique assemblages between Sickle, Pine, and Bear Creeks and through NMDS, Sickle Creek was most uniquely different with placement of brook trout and burbot, while Pine Creek had northern redbelly dace throughout. Pine Creek was also unique in the high abundance of brown and rainbow trout in the most downstream location (Huff Road), while Bear Creek was characterized by burbot only in downstream sections. Bear Creek also had longnose and western blacknose dace as unique contributors. In conclusion, the site-specific habitat template of each stream strongly dictated the fish species composition. For example, lower Bear Creek had much high concentrations of fine sediment and lower gradient – these two variables coupled with proximity to the Big Manistee River may have accounted for the high abundance of burbot found at this site. The habitat of lower Pine Creek consisted of relatively higher gradient and cooler water temperatures on account of the stream traversing a lateral

glacial moraine. These two factors coupled with the large abundance of large woody debris likely explain the high density of brown and rainbow trout. Further studies are underway to identify this relationship. Similarly, upstream Pine Creek was located at the outflow from a large and extensive marsh and therefore fine sediments were dominant and water temperatures were relatively warm.

Literature Cited:

- Allen, M., Combs, D.L., Cook, S.B., and M.R. Edwards. 2003. Comparison of single-pass electrofishing to depletion sampling for surveying fish assemblages in small warmwater streams. Journal of Freshwater Ecology. Vol 18, no. 4:625-634.
- Barbour, M.T., Gerritsen, J., Snyder, B.D., and J.B. Stribling. 1999. Rapid bioassessment protocols for use in wadeable streams and rivers. United States Environmental Protection Agency. 2nd Ed.
- Brazner, J.C., Tanner, D.K., Detenbeck, N.E., Batterman, S.L., Stark, S.L., Jagger, L.A., and V.M. Snarski. 2005. Regional, watershed, and site-specific environmental influences on fish assemblage structure and function in western Lake Superior tributaries. Canadian Journal of Fisheries and Aquatic Sciences, 62(6), 1254-1270
- Carline, R.F. 2001. Effects of high-frequency pulsed-DC electrofishing on a wild brown trout population. North American Journal of Fisheries Management: 21, 571-579.
- De Mol, N. 2007. Benthic macroinvertebrate response to road-stream crossing and stream bank improvements and longitudinal patterns in Bear, Pine, and Sickle Creeks, Manistee County, Michigan. MS Thesis, Grand Valley State University.
- Doonan, C.J., and G.E. Hendrickson. 1972. Reconnaissance of the Manistee River, a cold-water river in the northwestern part of Michigan's southern peninsula: U.S. Geological Survey Hydrologic Investigations Atlas 346, 2 sheets, scale 1:62,500.
- Grenouillet, G., Pont, D., and C. Herisse. 2004. Within-basin fish assemblage structure: the relative influence of habitat versus stream spatial position on local species richness. Canadian Journal of Fisheries and Aquatic Sciences: 61(1), 93-102
- Natural Resources Conservation Service. 2004. Landforms of Northern Lower Michigan: NRCS Michigan State OMCE.

- Page, L.M., and B.M. Burr. 1991. Freshwater Fishes. Houghton Mifflin Co., New York, New York.
- Paller, M. 1994. Relationships between Fish Assemblage Structure and Stream Order in South Carolina Coastal Plain Streams. Transactions of the American Fisheries Society: 123, 150–161
- Schaefer, J.F., and J.R. Kerfoot. 2004. Fish Assemblage Dynamics in an Adventitious Stream: A Landscape Perspective. American Midland Naturalist: 151(1), 134-145
- Smith, T.A., and C.E. Kraft. 2005. Stream Fish Assemblages in Relation to Landscape Position and Local Habitat Variables. Transactions of the American Fisheries Society: 134(1), 430-440
- Snyder, E.B., J. DeBoer, K. Nault. 2007. Biophysical response summaries in Bear, Pine, and Sickle Creeks, Manistee, MI. 2007. S. Ogren and J. Holtgren (co-principle investigators and editors) *In* Big Mainstee River Targeted Watershed Initiative Final Technical Report. Submitted to EPA. 74 pgs.
- Taylor, C.M., Holder, T.L., Fiorillo, R.A., Williams, L.R., Thomas, R.B., and M.L. Warren. 2006. Distribution, abundance, and diversity of stream fishes under variable environmental conditions. Canadian Journal of Fisheries and Aquatic Sciences, 63(1), 43-54
- Thomas, D.A. 2002. Fish and invertebrate community composition: a comparison of headwater and adventitious streams (Michigan). Masters Abstracts International: 40(4), 926
- United States Geological Survey. 1990. Land Cover Manistee County: Michigan State University NCCD, Michigan State University Board of Trustees.
- Wootton, R.J. 1998. Ecology of Teleost Fishes, 2nd Ed. Kluwer Academic Publishers, Boston, Massachusetts.

Tables:

Table 6. Analysis of variance results for various species CPUE for longitudinal distribution in Sickle Creek. Statistical analysis included repeated measures ANOVA. See table 1 for full names of location abbreviations.

Species	Locations compared and mean CPUE	p-value
Coho salmon	SIUS2 (0), SIDS3 (0.575)	0.078
	SIUS1 (0), SIDS3 (0.575)	0.078

Table 7. Analysis of variance results for various species CPUE for longitudinal distribution in Pine Creek. Statistical analysis included repeated measures ANOVA. See Table 1 for full names of location abbreviations.

Species	Locations and mean CPUE	p-value	Species	Locations and mean CPUE	p-value
W. blacknose dace	PLUS1 (2.67), PLDS1 (6.09)	0.070	Creek chub	PLDS3 (1.57), STDS3 (0.16)	0.011
	PLDS1 (6.09), Huff Road (0)	0.000		PLDS3 (1.57), Huff Road (0.01)	0.003
	PLDS2 (3.87), Huff Road (0)	0.023	Central mudminnow	PLUS2 (1.42), PLDS3 (0.32)	0.008
	PLDS3 (5.38), Huff Road (0)	0.000		PLUS2 (1.42), USFS 8430 (0.03)	0.033
	USFS 8430 (4.46), Huff Road (0)	0.005		PLUS2 (1.42), STDS1 (0.32)	0.002
	STUS1 (3.38), Huff Road (0)	0.075		PLUS2 (1.42), STDS3 (0.04)	0.033
	STDS1 (3.49), Huff Road (0)	0.058		PLUS2 (1.42), Huff Road (0.03)	0.003
Brown trout	PLUS2 (0.05), Huff Road (3.11)	0.000		PLUS1 (1.72), PLDS1 (0.63)	0.002
	PLUS1 (0). Huff Road (3.11)	0.000		PLUS1 (1.72), PLDS2 (0.16)	0.038
	PLDS1 (0.03), Huff Road (3.11)	0.000		PLUS1 (1.72), PLDS3 (0.32)	0.000
	PLDS2 (0.01), Huff Road (3.11)	0.000		PLUS1 (1.72), USFS 8430 (0.03)	0.002
	PLDS3 (0.02), Huff Road (3.11)	0.000		PLUS1 (1.72), STUS2 (0.46)	0.000
	USFS 8430 (0.42), Huff Road (3.11)	0.000		PLUS1 (1.72), STUS1 (0.40)	0.005
	STUS2 (0.21), Huff Road (3.11)	0.000		PLUS1 (1.72), STDS1 (0.32)	0.002
	STUS1 (0.16), Huff Road (3.11)	0.000		PLUS 1 (1.72), STDS2 (0.39)	0.004
	STDS1 (0.34), Huff Road (3.11)	0.000		PLUS1 (1.72), STDS3 (0.04)	0.000
	STDS2 (0.35), Huff Road (3.11)	0.000		PLUS1 (1.72), Huff Road (0.03)	0.000
	STDS3 (0.73), Huff Road (3.11)	0.000	N. redbelly dace	PLUS2 (3.78), STUS2 (0)	0.042
Creek chub	PLUS2 (2.03), PLDS2 (1.01)	0.008		PLUS2 (3.78), STDS2 (0)	0.042
	PLUS2 (2.03), PLDS3 (1.56)	0.033		PLUS2 (3.78), STDS3 (0.02)	0.043
	PLUS2 (2.03), USFS 8430 (0.43)	0.002		PLUS2 (3.78), Huff Road (0)	0.042
	PLUS2 (2.03), STDS1 (0.22)	0.033		PLDS1 (5.90), PLDS3 (1.71)	0.016
	PLUS2 (2.03), STDS3 (0.16)	0.003		PLDS1 (5.90), USFS 8430 (0.27)	0.000
	PLUS2 (2.03), Huff Road (0.01)	0.002		PLDS1 (5.90), STUS2 (0)	0.000
	PLUS1 (1.52), STUS2 (0.21)	0.023		PLDS1 (5.90), STUS1 (0.13)	0.000
	PLUS1 (1.52), STUS1 (0.07)	0.008		PLDS1 (5.90), STDS1 (0.10)	0.000
	PLUS1 (1.52), STDS1 (0.22)	0.024		PLDS1 (5.90), STDS2 (0)	0.000
	PLUS1 (1.52), STDS2 (0.20)	0.022		PLDS1 (5.90), STDS3 (0.02)	0.000
	PLUS1 (1.52), STDS3 (0.16)	0.016		PLDS1 (5.90), Huff Road (0)	0.000
	PLUS1 (1.52), Huff Road (0.01)	0.005	Rainbow trout	PLUS2 (0), Huff Road (1.87)	0.000
	PLDS1 (1.68), USFS 8430 (0.43)	0.033		PLUS1 (0.07), Huff Road (1.87)	0.000
	PLDS1 (1.68), STUS2 (0.21)	0.006		PLDS1 (0.30), Huff Road (1.87)	0.002
	PLDS1 (1.68), STUS1 (0.07)	0.002		PLDS2 (0.43), Huff Road (1.87)	0.005
	PLDS1 (1.68), STDS1 (0.22)	0.006		PLDS3 (0.29), Huff Road (1.87)	0.002
	PLDS1 (1.68), STDS2 (0.20)	0.006		USFS 8430 (0.45), Huff Road (1.87)	0.006
	PLDS1 (1.68), STDS3 (0.16)	0.004		STUS2 (0.24), Huff Road (1.87)	0.001
	PLDS1 (1.68), Huff Road (0.01)	0.004		STUS1 (0.31), Huff Road (1.87)	0.002
	PLDS3 (1.57), STUS2 (0.21)	0.016		STDS1 (0.40), Huff Road (1.87)	0.004
	PLDS3 (1.57), STUS1 (0.07)	0.005		STDS2 (0.63), Huff Road (1.87)	0.027
	PLDS3 (1.57), STDS1 (0.22)	0.017		STDS3 (0.67), Huff Road (1.87)	0.036
	PLDS3 (1.57), STDS2 (0.20)	0.015			
	PLDS3 (1.57), STDS3 (0.16)	0.011	1		

Table 8. Analysis of variance results for various species CPUE for longitudinal distribution in Bear Creek. Statistical analysis included repeated measures ANOVA.

Species	Locations and mean CPUE	p-value
		1
W. blacknose dace	SWAINDS (2.58), LOWER BEAR (0.08)	0.084
Burbot	MILKSUS (0.11), LOWER BEAR (1.83)	0.002
	MILKSDS (0.12), LOWER BEAR (1.83)	0.002
	SWAINDS (0.13), LOWER BEAR (1.83)	0.002
	SWAINDS2 (0.04), LOWER BEAR (1.83)	0.001
	JOHNSON (0.02), LOWER BEAR (1.83)	0.001
	SPIRITOFWOODS (0.29), LOWER BEAR (1.83)	0.006

Table 9. Fish species that differed significantly (ANOVA) between upper, mid, and lower reaches for Sickle, Pine, and Bear Creeks. Common species are also listed that indicate species that are fairly consistent along the longitudinal continuum.

	Sickle Creek (1st)	Pine Creek (2 nd)	Bear Creek (4 th)	Common Species
Upper	Brook trout	Northern redbelly dace Western blacknose dace	Western blacknose dace	Mottled sculpin Central mudminnow Chinook salmon Coho salmon
Middle		Northern redbelly dace Western blacknose dace	Longnose dace Western blacknose dace	Mottled sculpin Central mudminnow Chinook salmon Coho salmon
	•	-	•	•
Lower	Burbot	Brown trout Rainbow trout	Burbot	Mottled sculpin Central mudminnow Chinook salmon Coho salmon

Figures:




Figure 11. Sickle Creek longitudinal profile for Spring 2004 (A), Fall 2004 (B), Spring 2005 (C), and Fall 2005 (D). In Spring 2004, burbot and coho salmon increase from zero abundance above vs. below the perched culverts. Mottled sculpin were fairly consistent throughout and were the most dominant species in the entire stream. In Fall 2004, the same trends are seen with burbot increasing from up to down stream with brook trout only in upper reaches. In Spring 2005, mottled sculpin were the most dominant species throughout. Also, rainbow trout were highest above the culverts, while burbot, chinook salmon, and coho salmon were only found in downstream reaches. In Fall 2005, Mottled sculpin were the most dominant species throughout, while burbot and chinook salmon were only found in lower reaches. US = upstream, DS = downstream, culvert was located between US 1 and DS 1. The order of the legends matches the sequence of fish from top to bottom in the figure.




Figure 12. Pine Creek longitudinal profile for Spring 2004 (A), Fall 2004 (B), Spring 2005 (C), and Fall 2005 (D). In Spring 2004, brown and rainbow trout decreased in upper reaches but were dominant at Huff Road, the furthest reach downstream. Also, western blacknose dace were most dominant in mid-reaches and decreased in either direction, while northern redbelly dace were most abundant in headwater reaches. In Fall 2004, brown and rainbow trout increased from headwater reaches to downstream reaches. Western blacknose dace were most dominant at mid-reaches, while northern redbelly dace were most dominant in headwater reaches. In Spring 2005, chinook salmon abundance was higher in headwater reaches while creek chub were in low abundance in the lower reaches. Brown and rainbow trout were the most dominant at the Huff Road reach. In Fall 2005, northern redbelly dace were only located in the headwater reaches, while most brown and rainbow trout were located in downstream reaches. Western blacknose dace were dominant in most reaches, except for Huff Road.




Figure 13. Bear Creek longitudinal profile for Fall 2004 (A), Spring 2005 (B), and Fall 2005 (C). In Fall 2004, western blacknose dace increased in abundance from lower to upper reaches, while burbot were mainly located in lower reaches. Most other species were fairly consistent throughout the stream. In Spring 2005, western blacknose dace were in highest abundance in headwater reaches, while burbot were located only in downstream reaches. White sucker were in highest abundance in lower reaches. In Fall 2005, western blacknose dace and mottled sculpin were highest in upper reaches, while burbot and white sucker were highest at lower reaches.

Sickle Creek Fish Data




Figure 14. CPUE (fish/minute) data for various fish species in Sickle Creek. Reach abbreviations as in Table 1.

Pine Creek Fish Data




Figure 15. CPUE (fish/minute) data for various species in Pine Creek. Reach labels as in Table 1.

Bear Creek Fish Data




Figure 16. CPUE (fish/minute) data for various fish species in Bear Creek. Reach labels as in Table 1.




Figure 17. Discharge was positively correlated and explained 25.5% and 22.3% of the variation in species richness (A) and fish diversity (B) in Sickle Creek. pH was negatively correlated to fish density, but this was largely driven by a single data point and likely has limited ecological significance.





Figure 18. Discharge was negatively correlated with species richness (A) and species diversity (B) in Pine Creek.






Figure 19. Temperature was positively correlated with species dominance (A) while dissolved oxygen was negatively correlated with species diversity (B) in Bear Creek. Species diversity (C) was also negatively correlated with temperature in Bear Creek.




Figure 20. NMDS plot showing dispersal of fish assemblages at reaches and their relationship to water quality measurements in Sickle Creek. The blue triangles are the reaches with their respective name (as indicated in Table 1), and the red lines are the vectors for the joint bi-plot. The length of each vector is an indication of the strength of the relationship between that chemical parameter and the closest reach-level fish community. For example, both temperature and discharge are strongly associated with the fish communities at both SIDS1 and 2. Temp = temperature, DO = dissolved oxygen, pH = pH, SPC = specific conductivity, and Q = discharge.




Figure 21. NMDS plot showing dispersal of fish assemblages at reaches and their relationship towards water quality measurements in Pine Creek. The blue triangles are the reaches with their respective name (as indicated in Table 1), and the red lines are the vectors for the joint bi-plot. Temp = temperature, DO = dissolved oxygen, pH = pH, SPC = specific conductivity, and Q = discharge.




Figure 22. NMDS plot showing dispersal of fish assemblages at reaches and their relationship towards water quality measurements in Bear Creek. The blue triangles are the reaches with their respective name (as indicated in Table 1), and the red lines are the vectors for the joint bi-plot. The length of each vector shows how intense the relationship with the reaches are. Temp = temperature, pH = pH, and SPC = specific conductivity.

CHAPTER 4

FISH ASSEMBLAGE SPRING/SUMMER LONGITUDINAL PATTERNS IN THREE TRIBUTARIES OF THE LOWER MANISTEE RIVER, MICHIGAN

ABSTRACT

Fish migrate to various areas within and between coldwater watersheds, but few studies have examined the effects of seasonal differences with regard to entire stream, coldwater fish assemblages. The purpose of this study was to designate differences between whole fish assemblages and individual species among three tributaries of the lower Manistee River, Michigan, during spring and summer using a headwater to mouth approach in the two larger streams and a single location at the smallest stream. A total of 24 electrofishing reaches comprised the longitudinal gradient analysis. Whole fish assemblage measurements revealed no significant differences between seasons for Pine, Sickle, or Bear Creeks with respect to fish density, dominance, diversity, and richness. Sickle Creek (1st order) had no significant seasonal differences among any fish species present. In Pine Creek (2nd order), brook trout and coho salmon catch per unit effort were significantly higher in fall than spring. Bear Creek (4th order) had higher relative proportions of brook trout in spring versus fall samples. Most species from the streams appeared to remain in similar areas during these two seasons.

Introduction:

Fish migrate to various areas between and within coldwater watersheds and do so primarily to reach spawning areas, cooler water, and food resources. Diana et al. (2004) conducted telemetry evaluations of brown trout movements in the mainstream Au Sable River, Michigan and showed that brown trout remained near resting sites diurnally and moved various distances nocturnally. The brown trout returned to the same home resting sites as the previous year. Similarly, Curry et al. (2002) studied riverine brook trout populations with radio telemetry in the Kennebecasis River, New Brunswick, Canada and

found that while summer and winter movements were minimal, spring and fall movements were substantial—potentially for finding summer holding areas and spawning grounds. Trout species have been examined extensively because of their importance to the recreational fishing industry, but are only a small portion of the entire fish assemblage within a coldwater stream. Other fish species migrations are not as well known and documented.

Some studies have examined somewhat larger portions of stream fish assemblages. For instance, Magalhaes (1993) examined food resource use by seven cyprinids from an Iberian stream and found that differences in food resource use were found both between species and within species between seasons. In another study, Grossman et al. (1987), studied some members of a Mediterranean riverine fish assemblage in relation to microhabitat use. Most seasonal differences in microhabitat use were attributable to seasonal changes in microhabitat availability, although all species selectively occupied deeper microhabitats during spring and early summer.

Few studies have examined the effects of seasonal differences with regard to entire stream, coldwater fish assemblages. Meador and Matthews (1992) studied spatial and temporal patterns in fish assemblage structure of an intermittent Texas stream and found that despite drastic seasonal fluctuations in discharge, abundance of individual species varied more spatially among sites than temporally at individual sites. Jonsson (1991) showed that water flow, water temperature, and light are environmental variables that influence when fish migrate and the intensity of the migration itself and that these variables apply to both upstream and downstream migration, but their effects may vary among rivers and species.

While many studies have examined the effects of seasonality on movements of fish species, few have studied the entire fish assemblage at numerous locations, headwaters to mouth, and between three different order streams. The tributaries of the lower Manistee River offered an excellent opportunity to study the unique species associated with three different order streams and also to describe the seasonal differences seen with respect to not only individual species, but also the entire assemblages using various indices. The purpose of this study was to 1) designate differences between whole fish assemblages and individual species between spring and summer in three different order streams, and 2) hypothesize why seasonal changes in these different order streams may be variously impacted by sedimentation and serial disconnection.

Materials and Methods:

Seasonal Fish Assemblage Change Experimental design: Fish assemblages were monitored during spring and summer of 2004 and 2005 to determine change between seasons. Data were pooled for each season and was assessed along a longitudinal gradient and both entire fish assemblages and individual species were analyzed. Data were quantified using CPUE as in the previous chapters.

Total fish relative density, dominance, diversity, and richness were evaluated using repeated measures ANOVA for whole fish assemblages using seasonal fish assemblage data. Individual species relative density, catch per unit effort, and relative proportions were also evaluated using repeated measures ANOVA. Sickle Creek had 2 replicates for each season and 5 reaches with a total of 10 replicates per season. Pine Creek had 2 replicates with 12 reaches with a total of 24 reaches used per season. Bear

Creek had 1 replicate for spring sampling with 2 replicates for fall with a total of 7 and 14 for spring and fall, respectively. Each sample was considered a representative sample from each reach during each sampling period.

Results and Discussion:

Whole Fish Community Seasonal Differences: There were no significant differences found between seasons for Sickle, Pine, or Bear Creeks for density, dominance, diversity, and richness (Table 10, Figures 23-26). One possible reason why density, dominance, diversity, and richness between seasons for each stream was not significant was that if one species moved out, another species was there to replace it and if one species decreased in density, another species filled the space of the former. Many migratory species inhabit these streams at certain times of the year including rainbow trout, Chinook salmon, and coho salmon. These species may be migrating out and other more abundant species such as brown trout may displace those species when the other species do not seasonally inhabit the area. It appears that the potadromous species may not be significantly influencing the fish community in terms of relative density, dominance, diversity, and richness. More sampling in late fall and winter would yield more valuable insight into what is occurring in these streams.

Taxa-specific seasonal differences:

Pine Creek: Brook trout and coho salmon CPUE were found to be higher in summer than spring (Table 10 & 11, Figures 27-29). Coho salmon have been known to move out of rivers after the first year, although a few remain and could be considered residents. This

correlates with Welsh et al. (2001) who showed juvenile coho salmon reside in streams until smolting occurs, usually within two years at most. Seasonal migration may be a reason for higher CPUE for brook trout in summer versus spring, however, higher CPUE for coho salmon in summer may be the result of the smaller size of the fry earlier in the year. This species may have been freshly hatched and fairly untouched by electrofishing. Also, it may be that more individuals may have emigrated to the reaches electrofished in this study.

Higher CPUE for brook trout in summer could be a consequence of movements far upstream from other areas such as Lake Michigan, which is located roughly 20 miles downstream from Pine Creek, or the mainstream Big Manistee River, into which Pine Creek empties. It may be possible that lake brook trout may enter the smaller streams to spawn. Other reasons why brook trout had higher CPUE in summer include movement to spawn in headwaters, movement for water quality preferences, and fishing pressure. Petty et al. (2005) found that the spatial distribution of brook trout within a central Appalachian watershed was significantly correlated with spawning intensity and habitat features such as instream cover, stream depth and width, and riparian canopy cover; all variables that would change significantly from spring to summer, especially in response to the seasonal flow regime. Another study found that brook trout from the Sainte-Marguerite River, Quebec, Canada, first migrated downstream over a month in spring and adults then undertook upstream migration to spawning areas from July to September with larger individuals migrating earlier (Lenormand et al. 2004). Other reasons for higher brook trout CPUE in summer include movement to unsampled reaches in spring and inability to sample certain sections of reaches effectively due to higher water levels and resulting habitat.

Longnose dace density and CPUE were higher in spring versus summer. In fact, no longnose dace were found in this stream in summer. One reason could be the influence of larger predatory salmonid movement into the area. Another reason could be movement to reaches unsampled, but also the effect of many macroinvertebrates emerging as adults and dispersing out of the area. Though speculative, this compares to the findings of Thompson et al. (2001), which showed there was a significant, positive correlation between the biomass of benthic macroinvertebrates and longnose dace density in all seasons in a southern Appalachian stream. Perhaps various invertebrate taxa were not available in high abundances in summer versus spring.

Rainbow trout density and CPUE were higher in summer versus spring. Rainbow trout grow rapidly from hatching to the fall of the first year. Some rainbow trout in spring may have been overlooked from electrofishing size-selectivity, but also these species tend to move upstream to locate cooler water temperatures, food availability, and better-suited current velocities. Rainbow trout then migrate downstream usually after the second year, supporting the findings of very few larger rainbow trout within the streams. This compares with the findings of Daugherty et al. (2003) who showed rainbow trout had residence times of usually no more than two years in streams of the eastern Upper Peninsula of Michigan.

Other species present in Pine Creek had no significant seasonal differences. We hypothesized that brown trout would have large seasonal movements, but did not observe this pattern. Bettinger and Bettoli (2004) showed that the range of movement for brown

trout in the Clinch River, Tennessee, was significantly larger in fall than in any other season. Brown trout that were monitored for more than 1 year exhibited a limited range of movement during the winter, spring, and summer, but they made extensive movements during the fall season, presumably to spawn. In the current study, if seasonal movements of these individuals occurred, they may have happened later or earlier than when sampling was conducted.

As expected, another species not seasonally different was the Johnny darter; an observation supported by Mundahl and Ingersoll (1983) who found that only a small percentage of Johnny darters moved between seasons in an Ohio stream.

Sickle Creek: Sickle Creek had no seasonal differences among any fish species present, which was not surprising for species like the mottled sculpin. Downhower and Brown (1979) showed that even during spawning periods, mottled sculpin moved only small distances to find mates and that they are generally considered residents of the streams in which they reside.

Bear Creek: Brook trout had higher spring relative abundance than summer (Table 10 & 11). This could be a consequence of water quality preferences, fishing pressure, and the seasonal shift in abundances of other species. The change in migratory fish abundance may ultimately affect brook trout by displacing them to other areas of the stream, although both Chinook and coho salmon had higher spring density, CPUE, and relative proportions. Ultimately, it may be that in Pine Creek, that the increased summer coho salmon CPUE was the result of more individuals of this species remaining within the

stream reaches and that less individuals were within the stream reaches in spring. The same may be true for brook trout in both Pine and Bear Creeks.

The pattern in brook trout was originally hypothesized to be adversely affected by Chinook and coho salmon abundance. However, it appears that the potential effect of these two salmonids on brook trout may not be as adverse as previously thought. Brook trout had higher spring relative abundances in Bear Creek, along with both Chinook and coho salmon showing a similar trend. However, it may be that most brook trout were located in more upstream reaches, while Chinook and coho salmon were found in a more widespread distribution, although they decreased toward the furthest downstream reach. The brook trout in Bear Creek may have had a higher relative abundance in spring because they had moved to even further upstream reaches during the summer. Those reaches were not sampled as part of this study, although ongoing research has documented an evenly dominated community of brook and rainbow trout approximately 50 river kilometers upstream of the Milks Road site at the Leffew road stream crossing (Snyder et al. 2007). Alternatively, it may be that this pattern observed is not ecologically meaningful, and is simply due to chance.

Creek chub and mottled sculpin had higher summer CPUE values versus spring. Our data contradicts that of Pezold et al. (1997), who showed that in the Little Missouri River System in the Ouachita Mountains of Arkansas, creek chub appeared to be year-around residents with few seasonal movements. Downhower and Brown (1979), found that mottled sculpin are generally considered residents of the streams they reside in with little seasonal movement. We observed significant seasonal changes in sculpin CPUE in Bear Creek. In this system, temperature changes the most among the three streams

studied with values reaching 20°C in mid-summer. The increased creek chub and mottled sculpin CPUE in summer are likely to be from new arrivals of these species, either from emigration from the main channel or, more likely from emigration from unsampled adjacent reaches.

Sickle, Pine, and Bear Creeks each have visible problems associated with sedimentation and serial disconnection from improperly designed bridge and culvert placement. Little movement upstream of a perched culvert in Sickle Creek may have prohibited seasonal migration of some species. However, when flow was high enough to eliminate the plunge pool below the perched culvert, some species including trout that have physical attributes for swimming against high flow speeds, would potentially have been able to move upstream. Anthropogenic disturbances can affect migration patterns and management of tributaries of relatively large river systems is necessary to promote necessary seasonal movement patterns of the associated species.

Conclusion:

There were no significant differences found between seasons for Pine, Sickle, or Bear creeks for fish relative density, dominance, diversity, and richness. Seasonal fluctuations with individual species were mainly as expected. In Pine Creek, brook trout and coho salmon CPUE, and Rainbow trout density and CPUE were significantly higher in summer than spring. Sickle Creek had no significant seasonal differences among any fish species present. In Bear Creek, brook trout had significantly higher spring relative proportions than summer samples and both Chinook and coho salmon had significantly higher spring density, CPUE, and relative proportions. However, creek chub and mottled sculpin had significantly higher summer CPUE values than spring. Differences are likely

from physiological changes of habitat, physical, and biotic parameters needed for species fitness and survival. Most species from the streams appeared to remain in similar areas from season to season. However, these results suggest that regular sampling is necessary to establish long-term patterns and trends. Also, we recommend incorporation of more seasons into fish sampling to truly understand seasonal patterns. We also believe that management of tributary rivers is essential to the watershed scale maintenance of healthy and dynamic fish communities and that the results of studies such as this one can be used as an effective tool to better manage and conserve the fishery.

Literature Cited:

- Allen, M., Combs, D.L., Cook, S.B., and M.R. Edwards. 2003. Comparison of single-pass electrofishing to depletion sampling for surveying fish assemblages in small warmwater streams. Journal of Freshwater Ecology. Vol 18, no. 4:625-634.
- Barbour, M.T., Gerritsen, J., Snyder, B.D., and J.B. Stribling. 1999. Rapid bioassessment protocols for use in wadeable streams and rivers. United States Environmental Protection Agency. 2nd Ed.
- Bettinger, J.M., and P.W. Bettoli. 2004. Seasonal movement of brown trout in the Clinch River, Tennessee. North American Journal of Fisheries Management, 24(4):1480-1485
- Carline, R.F. 2001. Effects of high-frequency pulsed-DC electrofishing on a wild brown trout population. North American Journal of Fisheries Management: 21, 571-579.
- Curry, R.A., Sparks, D., and J. Van De Sande. 2002. Spatial and temporal movements of a riverine brook trout population. Transactions of the American Fisheries Society. 131:551-560.
- Daughtery, D.J., Sutton, T.M., and R.W. Greil. 2003. Life-history characteristics, population structure, and contribution of hatchery and wild steelhead in a Lake Huron tributary. Journal of Great Lakes Research: 29(3), 511-520.

- Diana, J.S., Hudson, J.P., and R.D. Clark, Jr. 2004. Movement patterns of large brown trout in the mainstream Au Sable River, Michigan. Transactions of the American Fisheries Society: 133, 34-44
- Downhower, J.F., and L. Brown. 1979. Seasonal changes in the social structure of a mottled sculpin (*Cottus bairdi*) population. Animal Behavior, 27:451-458
- Grossman, G.D., de Sostoa, A., Freeman, M.C., and J. Lobon-Cervia. 1987.

 Microhabitat use in a Mediterranean riverine fish assemblage. Oecologia: 73(4), 490-500
- Hubert, W.A., and C.C. Kohler. 1999. Inland Fisheries Management in North America. American Fisheries Society Bethesda, Maryland 2nd Ed: 167-191.
- Jonsson, N. 1991. Influence of Water Flow, Water Temperature and Light on Fish Migration in Rivers. Nordic Journal of Freshwater Research: 66, 20-35
- Lenormand, S., Dodson, J.J., and A. Menard. 2004. Seasonal and ontogenetic patterns in the migration of anadromous brook charr (*Salvelinus fontinalis*). Canadian Journal of Fisheries and Aquatic Sciences, 61(1): 54-67
- Magalhaes, M.F. 1993. Feeding of an Iberian stream cyprinid assemblage: seasonality of resource use in a highly variable environment. Oecologia: 96(2), 253-260
- Meador, M.R., and W.J. Matthews. 1992. Spatial and temporal patterns in fish assemblage structure of an intermittent Texas Stream. American Midland Naturalist: 127(1), 106-114
- Motulsky, H. 1995. Intuitive Biostatistics. Oxford University Press, New York: 255-258.
- Mundahl, N.D., and C.G. Ingersoll. 1983. Early autumn movements and densities of johnny (*Etheostoma nigrum*) and fantail (*E. flabellare*) darters in a southwestern Ohio stream. Ohio Journal of Science, 83(3): 103-108
- Petty, J.T., Lamonthe, P.J., and P.M. Mazik. 2005. Spatial and Seasonal dynamics of brook trout populations inhabiting a central Appalachian watershed. Transactions of the American Fisheries Society, 134(3): 572-587
- Pezold, F., Crump, B., and W. Flaherty. 1997. Seasonal patterns of fish abundance in two mountain creeks of the Little Missouri River drainage, Arkansas. Journal of Freshwater Ecology, 12(1): 51-60
- Seesholtz, A., Cavallo, B.J., Kindopp, J., and R. Kruth. 2004. Early life history of fishes in the San Francisco estuary and watershed. American Fisheries Society Symposium, 39:141-166

- Snyder, E.B., J. DeBoer, K. Nault. 2007. Biophysical response summaries in Bear, Pine, and Sickle Creeks, Manistee, MI. 2007. S. Ogren and J. Holtgren (co-principle investigators and editors) *In* Big Manistee River Targeted Watershed Initiative Final Technical Report. Submitted to EPA. 74 pgs
- Thompson, R.A., Petty, T.J., and D.G. Grossman. 2001. Multi-scale effects of resource patchiness on foraging behavior and habitat use by longnose dace, *Rhinichthys cataractae*. Freshwater Biology, 46(2): 145-160
- Welsh, H.H. Jr., Hodgson, G.R., Harvey, B.C., and M.F. Roche. 2001. Distribution of juvenile coho salmon in relation to water temperatures in tributaries of the Mattole River, California. North American Journal of Fisheries Management: 21, 464-470.

Tables:

Table 10. Mean density, CPUE, and relative abundance data for significant differences

between seasons. Standard deviations are in parenthesis.

	Mean density (ha)			Mean CPUE	are in pai		Mean relative abundance		
Species and location	spring	fall	p- value	spring	fall	p- value	spring	fall	
Brook trout (Bear Creek)							0.30 (0.20)	0.09 (0.03)	
Brook trout (Pine Creek)				0.05 (0.02)	0.16 (0.07)	0.026			
Chinook salmon (Bear Creek)	48 (2)	3 (2)	0.025	0.40 (0.19)	0.21 (0.12)	0.011	8.5 (0.77)	3.1 (0.44)	
Coho salmon (Bear Creek)	12 (6)	6 (2)	0.021	0.09 (0.04)	0.05 (0.02)	0.017	2.01 (0.78)	0.86 (0.35)	
Coho salmon (Pine Creek)				0.040 (0.02)	0.12 (0.05)	0.013			
Creek chub (Bear Creek)				0.07 (0.03)	0.39 (0.08)	0.021			
Longnose dace (Pine Creek)	25 (2.0)	0	0.038	0.10 (0.07)	0	0.034			
Mottled sculpin (Bear Creek)				0.10 (0.05)	0.46 (0.09)	0.049			
Rainbow trout (Pine Creek)	38 (31)	10. (4.0)	0.011	0.20 (0.08)	0.74 (0.25)	0.002			

Table 11. Summary of spring/summer fish community responses in Sickle, Pine, and Bear creeks. RA = relative abundance, n.s. = not significant at p > .05.

	Fish community metrics		Species-specific Responses							
Stream	Density, Richness, Diversity, Dominance	Chinook salmon	Coho salmon	Rainbow trout	Brook trout	Creek chub	Mottled sculpin			
Sickle	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.			
Pine Creek	n.s.	n.s.	CPUE (summer)	Density, CPUE (summer)	CPUE (summer)	n.s.	n.s.			
Bear Creek	n.s.	RA, CPUE, Density (spring)	RA, CPUE, Density (spring)	n.s.	RA (spring)	CPUE (summer)	CPUE (summer)			
	spring									
	summer									

Figures:




Figure 23. Relative seasonal fish density of Pine, Sickle, and Bear Creeks, Manistee County, Michigan. Each stream spring and fall from data are placed adjacent to each other respective of the stream and date. Pine, Sickle, and Bear Creek seasonal fish density are shown. Standard error bars are also given.




Figure 24. Seasonal fish dominance for Pine, Sickle, and Bear Creeks, Manistee County, Michigan. Dominance values closer to a value of 1 indicate that some species were much more abundant in those areas. These data values are the spring and fall values.




Figure 25. Seasonal species diversity of Pine, Sickle, and Bear Creeks, Manistee County, Michigan. Spring and fall data are shown in the figure. Standard error bars are also shown. The higher the value towards one, the more diverse the area. As the value reaches 0, the less diverse. Bear Creek has the highest diversity.




Figure 26. Seasonal fish species richness of Pine, Sickle, and Bear Creeks, Manistee County, Michigan. The number of species is what the species richness represents. Bear Creek has the highest species richness, with Sickle Creek having the lowest richness.




Figure 27. Relative fish density per hectare of all significant fish differences for each respective stream. The streams where the differences occur are shown above the respective species. All differences shown are p<.05.




Figure 28. Fish catch per minute of all significant fish differences for each respective stream. The streams where the differences occur are shown above the respective species. All differences shown are p<.05.




Figure 29. Fish relative abundance of all significant fish differences for Bear Creek. The streams where the differences occur are shown above the respective species. All differences shown are p<.05.

CHAPTER 5

OVERALL DISCUSSION

Effect of undersized road-stream crossings:

Sickle Creek had reduced diversity and increased dominance above a substantially perched culvert. In Pine and Bear creeks, fish assemblage response above and below impact sites was mixed. For example, undersized road-stream culverts reduced upstream habitat quality while eroding banks tended to reduce downstream habitat quality. Consequently, fish assemblage measurements were correlated with sediment composition at most sites, although this was not tested empirically in this thesis but is based on statistically significant differences in sediment composition above and below sediment sources (Holtgren and Ogren 2007). Most reaches above road-stream crossings had finer sediment buildup above culverts and larger substrate in downstream reaches, thus supporting our hypothesis that large amounts of finer substrate upstream would influence whole fish community metrics such as diversity and dominance. Moreover, downstream reaches generally had more diverse whole fish community values than those located above the road-stream crossing. Such trends suggest that undersized road-stream crossings that have been in place for long periods of time tend to accumulate sediment upstream--the amount which depends on the degree of constriction. Our results also suggest that when each road-stream crossing is restored, larger substrate and more diverse fish communities would exist above the road-stream crossing more similar to those below the crossing. Improvements to road-stream crossings should be done in such a way to maximize natural river structure and function.

Longitudinal fish distribution:

Few studies have employed a headwaters to mouth approach to Great Lakes Region stream fish distribution, impaired by high rates of sedimentation and barriers to fish migration (but see Seelbach and Wiley 1997, and Zorn et al. 1998). Results of our study showed that each stream had a distinctive fish community, usually attributed to a few select species as well as their longitudinal distribution. Sickle Creek had more brook trout upstream and more burbot in downstream reaches. Pine Creek had a large concentration of brown and rainbow trout downstream and more western blacknose and northern redbelly dace upstream. Bear Creek had more burbot downstream with more western blacknose and longnose dace upstream.

Most other studies have focused on fish longitudinal distribution and its relationship to biotic and abiotic factors (Zorn et al. 1998, Wang et al. 2003, Grenouillet et al. 2004, Schaefer and Kerfoot 2004, Torgersen et al. 2006, and many additional studies). In the Great Lakes region, specifically Michigan, few studies have been done. Zorn et al. (1998) examined fish distribution and abundance patterns within the lower peninsula and used low-flow yield and catchment area as independent variables. They determined that stream fishes responded in an individualistic manner to stream conditions, but also mentioned the need for additional species-specific studies. In the western United States, Torgersen et al. (2006) had compared gradients in fish assemblage structure among rivers and at multiple spatial scales and found spatial structuring of fish assemblages exhibited a generalized pattern of cold- and coolwater fish assemblage zones, but varied with temperature, especially in the warmest stream. Our study results

were similar with respect to cold- versus coolwater fish assemblages. Also, including sediment and a constricted/perched culvert factor, fish distribution patterns continued to follow similar patterns. Such results suggest that even though differences may be seen at smaller spatial scales (road-stream crossings, up- versus downstream comparisons), those differences do not affect the headwater to mouth continuum and may be considered a relatively small-scale problem unless the road-stream crossing is perched. That being said, there is a significant body of evidence that cumulative impacts will negatively affect stream ecosystems (Frissell et al. 1986, Bohn and Kershner 2002, Bond and Lake 2003), although less work has been done specifically on the cumulative effects of road-stream crossings per se (Harper and Quigley 2000, Wheeler et al. 2005). Once restoration activities commence, the smaller spatial scales differences can be alleviated and restore fish passage above and below these sites. Road-stream crossing and eroding banks seem to have a localized versus a watershed-scale impact and furthermore, funds for repairing the road-crossings were well spent and should have a beneficial affect on each stream.

In addition to this study, additional focus on whole fish communities and individual species, combined with a longitudinal approach, sedimentation, and fish passage between up- and downstream reaches, should support our findings. For instance, one smaller-scale study could focus on tracking individual species distribution within each stream. Larger-scale studies should focus on overall stream impacts from restoration activities from both road-stream crossing improvements and stream bank stabilizations. Time is an important consideration in these types of studies, as many of the species sampled may take years to establish a new dynamic equilibrium. Moreover, the study would be strengthened by additional years of post-restoration sampling

Stream restoration is very important tool for maintaining and improving stream ecosystem integrity and should be implemented at a much higher rate then it is today. Many studies have stressed the importance of stream restoration as a tool for dealing with problems of land use and for improving and enhancing fish migration, reproduction, forage, and successful completion of all life-history stages (Schlosser 1991, Allan and Flecker 1993, Harding et al. 1998, Ward et al. 2001, Jungwirth et al. 2002, Allan 2004, Bernhardt et al. 2005, Jansson et al. 2005, Palmer et al. 2005, Meyer et al. 2007, Jacobson and Galat 2008). In conclusion, Allan and Flecker (1993) note that the potential for recovery of damaged river ecosystems is considerable, and promoting restoration in these affected areas will prove to be beneficial for all organisms—including humans—that rely on lotic systems for goods and service, asthetics, and life in general.

Appendix A - Electrofishing Data

Sickle Creek Electrofishing Data

Sickle	Upstream	า #2
--------	----------	------

Spring 2004	7/7/2004			Summer 2004	8/10/2004		
Species	Number F	Proportion C	atch/Minute	Species	Number	Proportion	Catch/Minute
Mottled Sculpin	51	91.071	3.844	Brook Trout	1	2.326	0.095
Brook Trout	1	1.786	0.075	Brown Trout	2	4.651	0.190
Rainbow Trout	4	7.143	0.302	Mottled Sculpin	35	81.395	3.333
				Northern Brook Lamprey	2	4.651	0.190
				Rainbow Trout	3	6.977	0.286
Spring 2005	6/17/2005			Summer 2005	7/27/2005		
Species	Number F	Proportion C	atch/Minute	Species	Number	Proportion	Catch/Minute
Mottled Sculpin	26	89.655	2.500	Mottled Sculpin	24	100	2.88
Rainbow Trout	3	10.345	0.288	·			

Upstream	

Spring 2004	7/7/2004			Summer 2004	8/10/2004		
Species	Number F	Proportion Cat	ch/Minute	Species	Number	Proportion	Catch/Minute
Brook Trout	3	7.317	0.234	Brook Trout	4	11.429	0.501
Chinook Salmon	2	4.878	0.156	Brown Trout	2	5.714	0.251
Coho Salmon	2	4.878	0.156	Chinook Salmon	1	2.857	0.125
Mottled Sculpin	30	73.171	2.338	Coho Salmon	1	2.857	0.125
Northern Brook Lamprey	1	2.439	0.078	Creek Chub	2	5.714	0.251
Rainbow Trout	3	7.317	0.234	Mottled Sculpin	20	57.143	2.505
				Rainbow Trout	5	14.286	0.626
Spring 2005	6/17/2005			Summer 2005	7/27/2005		
Species	Number I	Proportion Cat	ch/Minute	Species	Number	Proportion	Catch/Minute
Mottled Sculpin	15	83.333	1.500	Brown Trout	1	6.250	0.131
Rainbow Trout	3	16.667	0.300	Mottled Sculpin	11	68.750	1.438
				Rainbow Trout	4	25.000	0.523

|--|

Spring 2004	6/30/2004			Summer 2004	8/9/2004		
Species		Proportion	Catch/Minute	Species	Number		Catch/Minute
Blacknose Dace	2	0.604	0.071	Brown Trout	1	0.847	0.074
Bluntnose Minnow	1	0.302	0.036	Burbot	27	22.881	2.010
Brook Stickleback	28	8.459	0.994	Chinook Salmon	4	3.390	0.298
Brook Trout	1	0.302	0.036	Creek Chub	7	5.932	0.521
Brown Trout	2	0.604	0.071	Johnny Darter	3	2.542	0.223
Burbot	28	8.459	0.994	Mottled Sculpin	74	62.712	5.509
Central Mudminnow	1	0.302	0.036	Northern Brook Lamprey	1	0.847	0.074
Chinook Salmon	6	1.813	0.213	Rainbow Trout	1	0.847	0.074
Coho Salmon	15	4.532	0.533				
Creek Chub	56	16.918	1.988				
Johnny Darter	2	0.604	0.071				
Mottled Sculpin	171	51.662	6.071				
Northern Brook Lamprey	3	0.906	0.107				
Northern Redbelly Dace	12	3.625	0.426				
Rainbow Trout	3	0.906	0.107				
Spring 2005	6/24/2005			Summer 2005	7/27/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Brown Trout	1	3.226	0.101	Burbot	3	5.769	0.357
Burbot	2	6.452	0.201	Chinook Salmon	2	3.846	0.238
Chinook Salmon	2	6.452	0.201	Johnny Darter	1	1.923	0.119
Coho Salmon	3	9.677	0.302	Mottled Sculpin	39	75.000	4.643
Johnny Darter	1	3.226	0.101	Northern Brook Lamprey	5	9.615	0.595
Mottled Sculpin	20	64.516	2.013	Rainbow Trout	2	3.846	0.238
Northern Redbelly Dace	1	3.226	0.101				
Rainbow Trout	1	3.226	0.101				

Sickle Downstream #2							
Spring 2004	6/23/2004			Summer 2004	8/9/2004		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Brown Trout	3	2.055	0.135	Brook Stickleback	1	1.493	0.098
Burbot	18	12.329	0.811	Brown Trout	1	1.493	0.098
Central Mudminnow	1	0.685	0.045	Burbot	25	37.313	2.451
Chinook Salmon	9	6.164	0.406	Chinook Salmon	1	1.493	0.098
Coho Salmon	14	9.589	0.631	Coho Salmon	3	4.478	0.294
Creek Chub	7	4.795	0.316	Johnny Darter	3	4.478	0.294
Johnny Darter	1	0.685	0.045	Mottled Sculpin	29	43.284	2.843
Mottled Sculpin	88	60.274	3.967	Northern Brook Lamprey	3	4.478	0.294
Northern Brook Lamprey	3	2.055	0.135	Rainbow Trout	1	1.493	0.098
Rainbow Trout	2	1.370	0.090				
Spring 2005	6/24/2005			Summer 2005	7/27/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Burbot	2	16.667	0.212	Burbot	2	10.000	0.240
Chinook Salmon	1	8.333	0.106	Chinook Salmon	1	5.000	0.120
Lamprey	1	8.333	0.106	Mottled Sculpin	15	75.000	1.804
Mottled Sculpin	8	66.667	0.847	Rainbow Trout	1	5.000	0.120
				White Sucker	1	5.000	0.120

Cialda Danmatus and #0							
Sickle Downstream #3	0/00/0004			0	0/0/0004		
Spring 2004	6/22/2004			Summer 2004	8/9/2004		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Burbot	6	13.043	0.615	Brown Trout	1	1.299	0.096
Chinook Salmon	2	4.348	0.205	Burbot	19	24.675	1.818
Coho Salmon	6	13.043	0.615	Central Mudminnow	1	1.299	0.096
Creek Chub	1	2.174	0.103	Chinook Salmon	1	1.299	0.096
Mottled Sculpin	28	60.870	2.872	Coho Salmon	5	6.494	0.478
Northern Brook Lamprey	1	2.174	0.103	Grass Pickerel	1	1.299	0.096
Rainbow Trout	2	4.348	0.205	Johnny Darter	3	3.896	0.287
				Mottled Sculpin	36	46.753	3.445
				Rainbow Trout	10	12.987	0.957
Spring 2005	6/23/2005			Summer 2005	7/27/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Burbot	4	6.667	0.401	Brown Trout	1	4.167	0.116
Central Mudminnow	1	1.667	0.100	Burbot	1	4.167	0.116
Chinook Salmon	1	1.667	0.100	Chinook Salmon	3	12.500	0.349
Coho Salmon	12	20.000	1.204	Creek Chub	1	4.167	0.116
Johnny Darter	1	1.667	0.100	Johnny Darter	6	25.000	0.698
Lamprey	1	1.667	0.100	Mottled Sculpin	8	33.333	0.930
Mottled Sculpin	38	63.333	3.813	Rainbow Trout	4	16.667	0.465
Rainbow Trout	2	3.333	0.201				

Pine Creek Electrofishing Data

Pine	Lake	Road	Upstream	#2
------	------	------	----------	----

7/16/2004			Summer 2004	8/11/2004		
Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
30	14.706	0.915	Blacknose Dace	50	26.60	3.61
5	2.451	0.153	Bluegill	6	3.19	0.43
2	0.980	0.061	Bluntnose Minnow	1	0.53	0.07
27	13.235	0.824	Brook Stickleback	3	1.60	0.22
1	0.490	0.031	Brook Trout	1	0.53	0.07
46	22.549	1.403	Brown Trout	1	0.53	0.07
18	8.824	0.549	Central Mudminnow	20	10.64	1.45
69	33.824	2.105	Creek Chub	35	18.62	2.53
2	0.980	0.061	Johnny Darter	1	0.53	0.07
4	1.961	0.122	Largemouth Bass	1	0.53	0.07
			Mottled Sculpin	1	0.53	0.07
			Northern Redbelly Dace	61	32.45	4.41
			White Sucker	7	3.72	0.51
7/11/2005			Summer 2005	9/16/2005		
	Proportion	Catch/Minuto			Proportion	Catch/Minuto
			•			4.537
						0.439
-						0.439
				-		0.585
						0.146
-						0.073
						1.976
_						2.854
						0.146
			,			0.140
2	1.250	0.122	<u> </u>	_		4.756
			,			0.073
			White Sucker	8	3.509	0.585
	Number 30 5 2 27 1 46 18 69 2 4	Number Proportion 30	Number Proportion Catch/Minute 30 14.706 0.915 5 2.451 0.153 2 0.980 0.061 27 13.235 0.824 1 0.490 0.031 46 22.549 1.403 18 8.824 0.549 69 33.824 2.105 2 0.980 0.061 4 1.961 0.122 7/11/2005 Number Proportion Catch/Minute 38 23.899 2.327 1 0.629 0.061 4 2.516 0.245 2 1.258 0.122 1 0.629 0.061 24 15.094 1.469 2 1.258 0.122 2 1.258 0.122 2 1.3836 1.347 63 39.623 3.857	Number	Number Proportion Catch/Minute Species Number 30 14.706 0.915 Blacknose Dace 50 5 2.451 0.153 Bluegill 6 2 0.980 0.061 Blunthose Minnow 1 27 13.235 0.824 Brook Stickleback 3 1 0.490 0.031 Brook Trout 1 46 22.549 1.403 Brown Trout 1 18 8.824 0.549 Central Mudminnow 20 69 33.824 2.105 Creek Chub 35 2 0.980 0.061 Johnny Darter 1 4 1.961 0.122 Largemouth Bass 1 Mottled Sculpin 1 Northern Redbelly Dace 61 White Sucker 7 7/11/2005 Summer 2005 8/16/2005 Number Species Number 8 23.899 2.327 Blacknose Dace 62	Number Proportion Catch/Minute Species Number Proportion 30 14.706 0.915 Blacknose Dace 50 26.60 5 2.451 0.153 Bluegill 6 3.19 2 0.980 0.061 Bluntnose Minnow 1 0.53 27 13.235 0.824 Brook Stickleback 3 1.60 1 0.490 0.031 Brook Trout 1 0.53 46 22.549 1.403 Brown Trout 1 0.53 18 8.824 0.549 Central Mudminnow 20 10.64 69 33.824 2.105 Creek Chub 35 18.62 2 0.980 0.061 Johnny Darter 1 0.53 4 1.961 0.122 Largemouth Bass 1 0.53 Northern Redbelly Dace 61 32.45 White Sucker 7 3.72 7/11/2005 Summer 2005 8/16/2005 Secies Numbe

Pine Lake Road Upstream #1

Spring 2004	7/14/2004			Summer 2004	8/11/2004		
Species	Number		Catch/Minute	Species	Number		Catch/Minute
Blacknose Dace	28	30.435	1.246	Blacknose Dace	53	31.93	4.08
Brook Stickleback	2	2.174	0.089	Bluegill	2	1.20	0.15
Central Mudminnow	6	6.522	0.267	Bluntnose Minnow	1	0.60	0.08
Creek Chub	20	21.739	0.890	Brook Stickleback	10	6.02	0.77
Longnose Dace	1	1.087	0.045	Brook Trout	4	2.41	0.31
Northern Redbelly Dace	32	34.783	1.424	Central Mudminnow	22	13.25	1.69
White Sucker	3	3.261	0.134	Creek Chub	30	18.07	2.31
				Northern Redbelly Dace	38	22.89	2.92
				Rainbow Trout	1	0.60	0.08
				White Sucker	5	3.01	0.38
Spring 2005	7/11/2005	;		Summer 2005	8/16/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	46	17.490	2.532	Blacknose Dace	38	26.027	2.850
Bluntnose Minnow	63	23.954	3.468	Bluegill	4	2.740	0.300
Brook Stickleback	7	2.662	0.385	Bluntnose Minnow	2	1.370	0.150
Central Mudminnow	45	17.110	2.477	Brook Stickleback	10	6.849	0.750
Common Shiner	1	0.380	0.055	Central Mudminnow	33	22.603	2.475
Creek Chub	21	7.985	1.156	Creek Chub	23	15.753	1.725
Northern Redbelly Dace	69	26.236	3.798	Northern Redbelly Dace	33	22.603	2.475
Rainbow Trout	1	0.380	0.055	Rainbow Trout	2	1.370	0.150
White Sucker	10	3.802	0.550	White Sucker	1	0.685	0.075

Spring 2004	eam #1 7/8/2004			Summer 2004	8/11/2004		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	141		6.957	Blacknose Dace	101	51.01	5.94
Bluegill	5	0.996	0.247	Bluegill	2	1.01	0.12
Brook Stickleback	2			Bluntnose Minnow	16		0.94
Brook Trout	1			Central Mudminnow	4		
Brown Trout	1			Creek Chub	17		1.00
Central Mudminnow	8			Largemouth Bass	1		0.06
Creek Chub	65			Mottled Sculpin	1		0.06
Longnose Dace	8			Northern Redbelly Dace	46		2.71
Mottled Sculpin	2			Rainbow Trout	7		
	260			White Sucker	1		0.41
Northern Redbelly Dace							
Rainbow Trout	6			Yellow Bullhead	2	1.01	0.12
White Sucker	3	0.598	0.148				
Spring 2005	7/7/2005		Ontale /Minute	Summer 2005	8/16/2005		Ontale /N Aliment
Species	Number		Catch/Minute	Species	Number		Catch/Minute
Blacknose Dace	112		6.575	Blacknose Dace	64		4.898
Bluegill	1			Bluegill	2		
Bluntnose Minnow	10			Bluntnose Minnow	12		
Central Mudminnow	22		1.292	Brook Trout	2		
Coho Salmon	3		0.176	Brown Trout	1		0.077
Creek Chub	29		1.703	Central Mudminnow	8		
Green Sunfish	3		0.176	Creek Chub	11		0.842
Mottled Sculpin	7			Johnny Darter	2		
Northern Redbelly Dace	84			Mottled Sculpin	5		0.383
Rainbow Trout	2		0.117	Northern Redbelly Dace	41		3.138
White Sucker	3	1.087	0.176	Rainbow Trout	5	3.205	0.383
				White Sucker	3	1.923	0.230
Pine Lake Road Downstr	eam #2						
				Summer 2004	8/12/2004		
	7/8/2004				6/12/2004		
Spring 2004	7/8/2004	Proportion	Catch/Minute				Catch/Minute
Spring 2004 Species	7/8/2004 Number		Catch/Minute	Species	Number	Proportion	Catch/Minute
Spring 2004 Species Blacknose Dace	7/8/2004 Number 77	40.104	3.717	Species Blacknose Dace	Number 71	Proportion 45.513	4.004
Spring 2004 Species Blacknose Dace Bluegill	7/8/2004 Number 77 6	40.104 3.125	3.717 0.290	Species Blacknose Dace Bluegill	Number 71	Proportion 45.513 2.564	4.004 0.226
Spring 2004 Species Blacknose Dace Bluegill Brook Trout	7/8/2004 Number 77 6 6	40.104 3.125 3.125	3.717 0.290 0.290	Species Blacknose Dace Bluegill Brook Trout	Number 71 4	Proportion 45.513 2.564 6.410	4.004 0.226 0.564
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner	7/8/2004 Number 77 6 6	40.104 3.125 3.125 0.521	3.717 0.290 0.290 0.048	Species Blacknose Dace Bluegill Brook Trout Brown Trout	Number 71 4 10	Proportion 45.513 2.564 6.410 0.641	4.004 0.226 0.564 0.056
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub	7/8/2004 Number 77 6 6 1 26	40.104 3.125 3.125 0.521 13.542	3.717 0.290 0.290 0.048 1.255	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow	Number 71 4 10 11 2	Proportion 45.513 2.564 6.410 0.641 2 1.282	3 4.004 4 0.226 0 0.564 0.056 2 0.113
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace	7/8/2004 Number 77 6 6 1 26 2	40.104 3.125 3.125 0.521 13.542 1.042	3.717 0.290 0.290 0.048 1.255 0.097	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub	Number 71 4 10 11 2 19 19	Proportion 45.513 4 2.564 6.410 0.641 1.282 12.179	4.004 4. 0.226 0.0564 0.056 2. 0.113 0.1071
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace	7/8/2004 Number 77 6 6 1 26 2 72	40.104 3.125 3.125 0.521 13.542 1.042 37.500	3.717 0.290 0.290 0.048 1.255 0.097 3.475	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin	Number 71 4 10 11 12 11 11 11 11 11 11 11 11 11 11 11	Proportion 45.513 2.564 6.410 0.641 1.282 12.179 0.641	4.004 0.226 0.564 0.056 0.113 0.1071 0.056
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout	7/8/2004 Number 77 6 6 1 26 2 72	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace	Number 71 4 10 11 12 11 11 11 11 11 11 11 11 11 11 11	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 12.179 0.641 0 25.641	4.004 0.226 0.564 0.056 0.113 0.1071 0.056 2.256
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout	7/8/2004 Number 77 6 6 1 26 2 72 1	40.104 3.125 3.125 0.521 13.542 1.042 37.500	3.717 0.290 0.290 0.048 1.255 0.097 3.475	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout	Number 71 4 10 11 12 11 11 11 11 11 11 11 11 11 11 11	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 1 12.179 0.641 0 25.641 3 5.128	4.004 0.226 0.0564 0.056 0.113 0.1071 0.056 2.256
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout	7/8/2004 Number 77 6 6 1 26 2 72 1 1	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout	Number 71 4 10 11 12 19 11 11 11 11 11 11 11 11 11 11 11 11	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 1 12.179 0.641 0 25.641 3 5.128	4.004 0.226 0.564 0.056 2.0113 1.071 0.056 2.256 0.451
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout	7/8/2004 Number 77 6 6 1 26 2 72 1	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout	Number 71 4 10 11 12 11 11 11 11 11 11 11 11 11 11 11	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 1 12.179 0.641 0 25.641 3 5.128	4.004 0.226 0.564 0.056 0.113 0.1071 0.056 2.256
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout	7/8/2004 Number 77 6 6 1 26 2 72 1 1	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout	Number 71 4 10 11 12 19 11 11 11 11 11 11 11 11 11 11 11 11	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 12.179 0.641 2 25.641 3 5.128 Proportion	4.004 0.226 0.0564 0.056 2.0113 0.1.071 0.056 2.256 0.451
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout Spring 2005 Species	7/8/2004 Number 77 6 6 1 26 2 72 1 1 7/7/2005 Number	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species	Number 71 4 10 11 12 11	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 0 12.179 0.641 0 25.641 3 5.128 6 Proportion 5 29.762	4.004 0.226 0.0.564 0.056 0.113 0.113 0.1.071 0.056 2.256 0.451 0.451
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout Spring 2005 Species Blacknose Dace	7/8/2004 Number 77 6 6 1 26 2 72 1 1 7/7/2005 Number 98	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048 Catch/Minute 5.833	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species Blacknose Dace	Number 71 4 10 11 12 12 12 12 12 12 12 12 12 12 12 12	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 7 12.179 0.641 0 25.641 3 5.128 6 Proportion 6 29.762 11.905	4.004 0.226 0.0.564 0.056 2.0.113 0.1.071 0.056 2.256 0.451 4. Catch/Minute 2.1.969 0.787
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout Spring 2005 Species Blacknose Dace Bluntnose Minnow	7/8/2004 Number 77 6 6 1 26 2 72 1 1 1 7/7/2005 Number 98 7	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495 3.535 0.505	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048 Catch/Minute 5.833 0.417	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species Blacknose Dace Brook Trout	Number 71 4 10 11 12 15 15 16 16 16 16 16 16 16 16 16 16 16 16 16	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 0 12.179 0.641 5 25.641 5 5.128 6 Proportion 6 29.762 11.905 2 2.381	4.004 0.226 0.564 0.056 2.0.113 0.1071 0.056 2.256 0.451 Catch/Minute 1.969 0.787 0.157
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Spring 2005 Species Blacknose Dace Bluntnose Minnow Brook Trout Central Mudminnow	7/8/2004 Number 77 6 6 1 26 2 72 1 1 1 7/7/2005 Number 98 7	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495 3.535 0.505 3.030	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048 Catch/Minute 5.833 0.417 0.060 0.357	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species Blacknose Dace Brook Trout Central Mudminnow Creek Chub	Number 71 4 10 11 12 15 15 16 16 17 18 18 18 18 18 18 18 18 18 18 18 18 18	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 0 12.179 0.641 5 25.641 5 128 6 6 6 7 7 8 7 8 9 7 8 9 10.714	4.004 4.0.226 0.564 0.056 2.0.113 0.1.071 0.056 2.256 0.451 1.021 1.0
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout Spring 2005 Species Blacknose Dace Bluntnose Minnow Brook Trout Central Mudminnow Coho Salmon	7/8/2004 Number 77 6 6 1 26 2 72 1 1 7/7/2005 Number 98 7 1 6 6	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495 3.535 0.505 3.030 0.505	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048 Catch/Minute 5.833 0.417 0.060 0.357	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species Blacknose Dace Brook Trout Central Mudminnow Creek Chub Mottled Sculpin	Number 71 4 10 11 12 11	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 12.179 0.641 0 25.641 3 5.128 Proportion 6 29.762 11.905 2 2.381 10.714 3 .571	4.004 4.0.226 0.564 0.056 2.0.113 0.1.071 0.056 2.256 0.451 Catch/Minute 2.1.969 6.0.787 0.157 0.709 0.236
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout Spring 2005 Species Blacknose Dace Bluntnose Minnow Brook Trout Central Mudminnow Coho Salmon Creek Chub	7/8/2004 Number 77 6 6 1 26 2 72 1 1 7/7/2005 Number 98 7 1 6 1 1	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495 3.535 0.505 3.030 0.505 8.586	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048 Catch/Minute 5.833 0.417 0.060 0.357 0.060 1.012	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species Blacknose Dace Brook Trout Central Mudminnow Creek Chub Mottled Sculpin	Number 71 4 10 10 11 12	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 12.179 0.641 0 25.641 6 5.128 Proportion 6 29.762 11.905 2 2.381 1 0.714 6 3.571 2 9.762	4.004 4.0.226 0.0.564 0.056 2.0.113 1.071 0.056 2.256 0.451 Catch/Minute 2.1.969 0.787 0.157 0.709 0.236 2.1969 1.969
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout Spring 2005 Species Blacknose Dace Bluntnose Minnow Brook Trout Central Mudminnow Coho Salmon Creek Chub Green Sunfish	7/8/2004 Number 77 6 6 1 26 2 72 1 1 7/7/2005 Number 98 7 1 6 1 1 177	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495 3.535 0.505 3.030 0.505 8.586	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048 Catch/Minute 5.833 0.417 0.060 0.357 0.060	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species Blacknose Dace Brook Trout Central Mudminnow Creek Chub Mottled Sculpin	Number 71 4 10 11 12 11	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 12.179 0.641 0 25.641 6 5.128 Proportion 6 29.762 11.905 2 2.381 1 0.714 6 3.571 2 9.762	4.004 4.0.226 0.0.564 0.056 2.0.113 1.071 0.056 2.256 0.451 Catch/Minute 2.1.969 0.787 0.157 0.709 0.236 2.1969 1.969
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout Spring 2005 Species Blacknose Dace Bluntnose Minnow Brook Trout Central Mudminnow Coho Salmon Creek Chub Green Sunfish Mottled Sculpin	7/8/2004 Number 77 6 6 1 26 2 72 1 1 7/7/2005 Number 98 7 1 6 1 1 17	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495 3.535 0.505 3.030 0.505 8.586 0.505	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048 Catch/Minute 5.833 0.417 0.060 0.357 0.060 1.012 0.060 0.060	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species Blacknose Dace Brook Trout Central Mudminnow Creek Chub Mottled Sculpin	Number 71 4 10 10 11 12	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 12.179 0.641 0 25.641 6 5.128 Proportion 6 29.762 11.905 2 2.381 1 0.714 6 3.571 2 9.762	4.004 4.0.226 0.0.564 0.056 2.0.113 1.071 0.056 2.256 0.451 Catch/Minute 2.1.969 0.787 0.157 0.709 0.236 2.1969 1.969
Spring 2004 Species Blacknose Dace Bluegill Brook Trout Common Shiner Creek Chub Longnose Dace Northern Redbelly Dace Rainbow Trout Rainbow Trout Spring 2005 Species Blacknose Dace Bluntnose Minnow Brook Trout Central Mudminnow Coho Salmon Creek Chub Green Sunfish	7/8/2004 Number 77 6 6 1 26 2 72 1 1 7/7/2005 Number 98 7 1 6 1 1 177	40.104 3.125 3.125 0.521 13.542 1.042 37.500 0.521 0.521 Proportion 49.495 3.535 0.505 3.030 0.505 8.586 0.505 0.505 29.798	3.717 0.290 0.290 0.048 1.255 0.097 3.475 0.048 0.048 Catch/Minute 5.833 0.417 0.060 0.357 0.060	Species Blacknose Dace Bluegill Brook Trout Brown Trout Central Mudminnow Creek Chub Mottled Sculpin Northern Redbelly Dace Rainbow Trout Summer 2005 Species Blacknose Dace Brook Trout Central Mudminnow Creek Chub Mottled Sculpin	Number 71 4 10 10 11 12	Proportion 45.513 4 2.564 6.410 0.641 2 1.282 12.179 0.641 0 25.641 6 5.128 Proportion 6 29.762 11.905 2 2.381 1 0.714 6 3.571 2 9.762	4.004 4.0.226 0.0.564 0.056 2.0.113 1.071 0.056 2.256 0.451 Catch/Minute 2.1.969 0.787 0.157 0.709 0.236 2.1969 1.969

Spring 2004	7/13/2004			Summer 2004	8/12/2004		
Species	Number Pi	roportion Cat	ch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	145	52.920	5.894	Blacknose Dace	126	57.014	6.29
Bluegill	4	1.460	0.163	Bluegill	2	0.905	0.100
Bluntnose Minnow	1	0.365	0.041	Brook Trout	4	1.810	0.200
Brook Trout	4	1.460	0.163	Central Mudminnow	1	0.452	0.050
Central Mudminnow	7	2.555	0.285	Coho Salmon	2	0.905	0.100
Common Shiner	1	0.365	0.041	Creek Chub	42	19.005	2.09
Creek Chub	21	7.664	0.854	Mottled Sculpin	4	1.810	0.200
Longnose Dace	4	1.460	0.163	Northern Redbelly Dace	e 30	13.575	1.49
Mottled Sculpin	1	0.365	0.041	Rainbow Trout	10	4.525	0.499
Northern Redbelly Dace	85	31.022	3.455				
White Sucker	1	0.365	0.041				
Spring 2005	7/7/2005			Summer 2005	8/16/2005		
Species	Number Pi	roportion Cat	ch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	94	63.514	5.562	Blacknose Dace	49	38.889	3.77
Bluntnose Minnow	2	1.351	0.118	Bluegill	4		0.30
Brook Stickleback	1	0.676	0.059	Brook Trout	8		0.616
Brook Trout	1	0.676	0.059	Central Mudminnow	7	5.556	0.539
Brown Trout	1	0.676	0.059	Creek Chub	27	21.429	2.08
Central Mudminnow	7	4.730	0.414	Mottled Sculpin	9		0.693
Coho Salmon	1	0.676	0.059	Northern Redbelly Dace		11.111	1.07
Creek Chub	21	14.189	1.243	Rainbow Trout	7		0.539
Mottled Sculpin	3	2.027	0.178	Yellow Bullhead	1	0.794	0.07
	14	9.459	0.828	Tellow Bullicad		0.754	0.07
•							
Rainbow Trout White Sucker	2	1.351 0.676	0.118 0.059				
Rainbow Trout White Sucker USFS 8430	2	1.351 0.676	0.118	Summer 2004	8/11/2004		
Rainbow Trout White Sucker USFS 8430 Spring 2004	2 1 7/14/2004 Number	1.351 0.676 Proportion	0.118	Summer 2004 Species Num			Catch/Minute
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species	7/14/2004	1.351 0.676 Proportion 0.935	0.118 0.059			Proportion (65.517	
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill	2 1 7/14/2004 Number	1.351 0.676 Proportion 0.935	0.118 0.059 Catch/Minute	Species Num	nber I		5.556
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace	7/14/2004 Number	1.351 0.676 Proportion 0.935 5 52.336	0.118 0.059 Catch/Minute 0.048	Species Num Blacknose Dace	nber I 114	65.517	5.556 0.049
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout	7/14/2002 Number	1.351 0.676 4 Proportion 0.935 6 52.336 0 9.346	0.118 0.059 Catch/Minute 0.048 2.710	Species Num Blacknose Dace Brook Trout	nber I 114 1	65.517 0.575	5.556 0.049 0.536
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout	7/14/200 ² Number	1.351 0.676 Proportion 0.935 5 52.336 0 9.346 0.935	0.118 0.059 Catch/Minute 0.048 2.710 0.484	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub	nber I 114 1 11	65.517 0.575 6.322	5.556 0.049 0.536 0.439
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub	7/14/2002 Number	1.351 0.676 Proportion 0.935 5 52.336 0 9.346 0.935 4 3.738	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter	nber I 114 1 11 9	65.517 0.575 6.322 5.172 0.575	5.556 0.049 0.536 0.439 0.049
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter	7/14/2004 Number 10 10	1.351 0.676 Proportion 0.935 5.2.336 9.346 0.935 4.3.738 0.935	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass	nber I 114 1 11 9 1	65.517 0.575 6.322 5.172 0.575 0.575	5.556 0.049 0.536 0.439 0.049
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace	7/14/2004 Number 156	1.351 0.676 Proportion 0.935 5.2.336 0.935 4.3.738 0.935 8.2.804	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.194	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin	nber I 114 1 11 9 1 1	65.517 0.575 6.322 5.172 0.575 0.575 8.046	5.556 0.049 0.536 0.439 0.049 0.049
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin	7/14/2004 Number	1.351 0.676 Proportion 0.935 5.52.336 0.935 4.0.935 4.0.935 8.2.804 8.2.804	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout	nber I 114 1 11 9 1 1 14	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471	5.556 0.049 0.536 0.439 0.049 0.049 0.682
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow	7/14/200 ² Number 1 56 10 1 2 2 3 3 3 4 2	1.351 0.676 Proportion 0.935 5.52.336 0.935 4.0.935 4.0.935 8.2.804 8.2.804 4.3.738	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin	nber I 114 1 11 9 1 1	65.517 0.575 6.322 5.172 0.575 0.575 8.046	5.556 0.049 0.536 0.439 0.049 0.682 0.634
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace	7/14/2004 Number	1.351 0.676 Proportion 0.935 5 52.336 0 9.346 0.935 4 3.738 0.935 8 2.804 4 3.738 0 18.692	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout	nber I 114 1 11 9 1 1 14	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471	Catch/Minute 5.556 0.049 0.536 0.439 0.049 0.682 0.633 0.487
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker	7/14/2002 Number 1 56 10 11 2 2 20 20 20 20 20 20 20 20 20 20 20 20	1.351 0.676 Proportion 0.935 5.52.336 0.935 4.3.738 0.935 8.2.804 8.2.804 4.3.738 9.18.692 4.3.738	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.194 0.968	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout	nber I 114 1 11 9 1 1 14	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471	5.556 0.049 0.536 0.439 0.049 0.049 0.682
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005	7/14/2004 Number 156 10 14 24 20 20	1.351 0.676 Proportion 0.935 5.2.336 0.935 4.3.738 0.935 8.2.804 8.2.804 4.3.738 0.18692 4.3.738	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.145 0.194	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker	114 114 11 11 9 1 1 14 13 10	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747	5.556 0.049 0.536 0.439 0.049 0.682 0.634 0.483
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species	7/14/2002 Number 1 56 10 1 4 2 20 2 7/6/2005 Number	1.351 0.676 Proportion 0.935 52.336 0.935 4 0.935 4 0.935 8 2.804 8 2.804 4 3.738 0 18.692 4 3.738 0 Proportion	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.145 0.194	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num	114 11 11 11 9 11 14 13 10	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747	5.556 0.049 0.534 0.433 0.044 0.683 0.634 0.483
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace	7/14/2002 Number 156 10 12 22 20 27/6/2005 Number 111	1.351 0.676 Proportion 0.935 5.52.336 0.935 4.3.738 1.0.935 8.2.804 4.3.738 0.18.692 4.3.738 0.738 0.738 0.738 0.738	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.045 0.145 0.145 0.194 0.968 0.194	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace	114 11 11 11 9 1 1 14 13 10 8/10/2005	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747	5.556 0.049 0.531 0.439 0.049 0.683 0.634 0.487
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace Brown Trout	7/14/2002 Number 156 10 11 22 20 27/6/2005 Number	1.351 0.676 Proportion 0.935 5 52.336 0 9.346 0.935 4 3.738 8 2.804 4 3.738 0 18.692 4 3.738 0 Proportion 72.549 4 2.614	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.145 0.194 0.968 0.194 Catch/Minute 5.925 0.214	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace Brook Trout	8/10/2005 bber I 114 11 11 12 14 13 10	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747	5.55 0.04 0.53 0.43 0.04 0.68 0.63 0.48 Catch/Minute 3.65 0.12
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace Brown Trout Brook Trout	7/14/2004 Number 156 10 11 2 20 7/6/2005 Number 111 2 2 2	1.351 0.676 Proportion 0.935 5.2.336 0.935 4.3.738 0.935 8.2.804 4.3.738 0.18.692 4.3.738 Proportion 72.549 4.2.614 2.2.614	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.194 0.968 0.194 Catch/Minute 5.925 0.214 0.107	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace Brook Trout Brown Trout	8/10/2005 8/10/2005 10 8/10/2005 10 10	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747	5.55i 0.04: 0.53i 0.43: 0.04: 0.68: 0.63: 0.48' Catch/Minute 3.65: 0.12: 0.12:
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace Brown Trout Brook Trout Central Mudminnow	7/14/2004 Number 156 10 11 22 20 7/6/2005 Number 111 22 22 23 24 25 26 26 27 27 27 27 27 27 27 27 27 27 27 27 27	1.351 0.676 Proportion 0.935 5.52.336 0.935 4.3.738 0.935 8.2.804 4.3.738 1.692 1.3738 Proportion 72.549 4.2.614 2.614 2.1.307	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.194 0.968 0.194 Catch/Minute 5.925 0.214 0.107	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub	8/10/2005 bber I 114 11 9 1 1 14 13 10 8/10/2005 bber I 60 2 7	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747 Proportion 0 50.420 1.681 5.882 8.403	5.55 0.04 0.53 0.43 0.04 0.68 0.63 0.48 Catch/Minute 3.65 0.12 0.42
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace Brown Trout Brook Trout Central Mudminnow Creek Chub	7/14/2004 Number 156 10 11 24 20 7/6/2005 Number 111 24 25	1.351 0.676 Proportion 0.935 5.2.336 9.346 0.935 8.2.804 8.2.804 8.3.738 9.18.692 9.3.738 9.18.692 9.2.614 1.307 2.614 2.1.307 2.1.307 5.882	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.194 0.968 0.194 Catch/Minute 5.925 0.214 0.107 0.107	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter	8/10/2005 iber I 8/10/2005 iber I 60 2 7 10 2	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747	5.55i 0.04i 0.53i 0.43i 0.04i 0.68i 0.63i 0.48i Catch/Minute 3.65i 0.12i 0.42i 0.61i
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace Brown Trout Brook Trout Central Mudminnow Creek Chub Johnny Darter	7/14/2002 Number 1 56 10 1 2 2 20 7/6/2005 Number 111 2 2 2 2 2 2 2 3 3 3 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	1.351 0.676 Proportion 0.935 5.52.336 0.935 4.3.738 3.2.804 4.3.738 18.692 4.3.738 Proportion 72.549 4.2.614 2.1.307 2.1.307	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.194 0.968 0.194 Catch/Minute 5.925 0.214 0.107 0.480 0.107	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass	8/10/2005 aber I 8/10/2005 bber I 10 2 1	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747 Proportion 0 50.420 1.681 5.882 8.403 1.681 0.840	5.55i 0.04i 0.53i 0.43i 0.04i 0.68i 0.63i 0.48i Catch/Minute 3.65i 0.12i 0.42i 0.61i 0.12i
Northern Redbelly Dace Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Greek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker	7/14/2002 Number 156 10 12 20 20 7/6/2005 Number 111 22 23 24 25 26 26 27 28	1.351 0.676 Proportion 0.935 5.52.336 0.935 4.3.738 3.2.804 4.3.738 0.935 3.2.804 4.3.738 0.935 6.2.804 4.3.738 0.935 6.2.804 4.3.738 0.935 18.692 4.3.738 0.935 18.692 1.307 2.614	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.194 0.968 0.194 Catch/Minute 5.925 0.214 0.107 0.480 0.107	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin	8/10/2005 abber 8/10/2005 bber 60 2 7 10 2 1 14	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747 Proportion (50.420 1.681 5.882 8.403 1.681 0.840 11.765	5.556 0.048 0.538 0.048 0.048 0.682 0.634 0.487 Catch/Minute 3.658 0.122 0.427 0.610 0.122
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace Brown Trout Brook Trout Central Mudminnow Orther Spring 2005 Species Company Darter Central Mudminnow Creek Chub Johnny Darter Mottled Sculpin Northern Redbelly Dace	7/14/2004 Number 10 11 2 2 7/6/2005 Number 111 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	1.351 0.676 Proportion 0.935 5.52.336 0.935 4.3.738 0.935 8.2.804 8.2.804 8.3.738 9.18.692 1.3.738 1.3	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.194 0.968 0.194 Catch/Minute 5.925 0.214 0.107 0.107 0.480 0.107	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout	8/10/2005 sheer I I I I I I I I I I I I I I I I I I	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747 Proportion (50.420 1.681 5.882 8.403 1.681 0.840 0.840 11.765	5.556 0.048 0.534 0.433 0.044 0.683 0.483 0.483 0.483 0.421 0.612 0.422 0.610 0.122 0.066 0.854
Rainbow Trout White Sucker USFS 8430 Spring 2004 Species Bluegill Blacknose Dace Brown Trout Brook Trout Creek Chub Johnny Darter Longnose Dace Mottled Sculpin Central Mudminnow Northern Redbelly Dace White Sucker Spring 2005 Species Blacknose Dace Brown Trout Brook Trout Central Mudminnow Cerek Chub Johnny Darter Modified Sculpin	7/14/2002 Number 156 10 12 20 20 7/6/2005 Number 111 22 23 24 25 26 26 27 28	1.351 0.676 Proportion 0.935 5.2.336 0.935 3.738 0.935 8.2.804 4.3.738 0.935 9.346 0.935 9.346 0.935 1.307 1.307 2.614 2.614 2.614 2.614 2.614 2.614 2.614 2.614 2.614 2.614 2.614 2.614 2.614 3.738	0.118 0.059 Catch/Minute 0.048 2.710 0.484 0.048 0.194 0.048 0.145 0.145 0.194 0.968 0.194 Catch/Minute 5.925 0.214 0.107 0.480 0.107	Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin Rainbow Trout White Sucker Summer 2005 Species Num Blacknose Dace Brook Trout Brown Trout Creek Chub Johnny Darter Largemouth Bass Mottled Sculpin	8/10/2005 abber I 8/10/2005 bber I 10 2 11 14	65.517 0.575 6.322 5.172 0.575 0.575 8.046 7.471 5.747 Proportion (50.420 1.681 5.882 8.403 1.681 0.840 11.765	5.556 0.048 0.536 0.048 0.048 0.682 0.634 0.487 Catch/Minute 3.658 0.122 0.427 0.611

Spring 2004	7/15/2004			Summer 2004	8/13/2004		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	72	78.261	3.303	Blacknose Dace	36	52.174	3.956
Bluegill	1	1.087	0.046	Bluegill	1	1.449	0.110
Brook Trout	1	1.087	0.046	Brown Trout	4	5.797	0.440
Brown Trout	2	2.174	0.092	Central Mudminnow	12	17.391	1.319
Coho Salmon	1	1.087	0.046	Coho Salmon	1	1.449	0.110
Creek Chub	3	3.261	0.138	Creek Chub	5	7.246	0.549
Mottled Sculpin	9	9.783	0.413	Mottled Sculpin	2	2.899	0.220
Rainbow Trout	2	2.174	0.092	Rainbow Trout	5	7.246	0.549
White Sucker	1	1.087	0.046	White Sucker	3		
Spring 2005	7/6/2005			Summer 2005	8/10/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	41	57.746	2.228	Blacknose Dace	36	58.065	2.842
Brook Trout	1	1.408	0.054	Brook Trout	3	4.839	0.237
Brown Trout	3	4.225	0.163	Brown Trout	2	3.226	0.158
Central Mudminnow	1	1.408	0.054	Central Mudminnow	6	9.677	0.474
Creek Chub	3	4.225	0.163	Coho Salmon	2	3.226	0.158
Johnny Darter	2	2.817	0.109	Mottled Sculpin	8	12.903	0.632
Mottled Sculpin	14	19.718	0.761	Rainbow Trout	2	3.226	0.158
Rainbow Trout	3	4.225	0.163	White Sucker	3	4.839	0.237
Red ear Sunfish	1	1.408	0.054				
White Sucker	2	2.817	0.109				
Steinberg Road Upstro	eam #1						
Spring 2004	7/14/2004	4		Summer 2004	8/13/2004		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	44	4 73.333	3 2.505	Blacknose Dace	84	65.116	6.139
Brown Trout	2	2 3.333	0.114	Brown Trout	3	2.326	0.219
Central Mudminnow	2	2 3.333	0.114	Central Mudminnow	10	7.752	0.731
Creek Chub		1 1.667	0.057	Chinook Salmon	1	0.775	0.073
Johnny Darter		1 1.667	0.057	Coho Salmon	2	1.550	0.146
Mottled Sculpin		6 10.000		Creek Chub	4		0.292
Rainbow Trout	;	3 5.000		Mottled Sculpin	12		
White Sucker	•	1 1.667	0.057	Northern Redbelly Dace			0.292
				Rainbow Trout	7		
				White Sucker	2	1.550	0.146
Carina 200E	7/0/000	-		Cummar 200F	0/40/0005	•	

			Rainbow Trout	7	5.426	0.512
						0.012
			White Sucker	2	1.550	0.146
/2005	i		Summer 2005	8/10/2005		
ber	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
51	53.125	3.048	Blacknose Dace	23	42.593	1.865
1	1.042	0.060	Brook Stickleback	1	1.852	0.081
6	6.250	0.359	Brown Trout	4	7.407	0.324
2	2.083	0.120	Central Mudminnow	5	9.259	0.405
4	4.167	0.239	Coho Salmon	3	5.556	0.243
2	2.083	0.120	Mottled Sculpin	13	24.074	1.054
18	18.750	1.076	Rainbow Trout	3	5.556	0.243
9	9.375	0.538	White Sucker	2	3.704	0.162
3	3.125	0.179				
	51 51 6 2 4 2 18	51 53.125 1 1.042 6 6.250 2 2.083 4 4.167 2 2.083 18 18.750 9 9.375	Description Description	Summer 2005 Summer 2005 Species Species Species Sp	Summer 2005 Summer 2005 Summer 2005 Summer 2005 Summer 2005 Number	Summer 2005 Summer 2005

Steinberg Road Downstr	eam #1						
Spring 2004	7/15/2004			Summer 2004	8/16/2004		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	45	71.429	2.290	Blacknose Dace	64	62.136	4.689
Brown Trout	5	7.937	0.254	Bluegill	3	2.913	0.220
Central Mudminnow	1	1.587	0.051	Brown Trout	9	8.738	0.659
Creek Chub	1	1.587	0.051	Central Mudminnow	2	1.942	0.147
Mottled Sculpin	6	9.524	0.305	Coho Salmon	2	1.942	0.147
Rainbow Trout	4	6.349	0.204	Johnny Darter	2	1.942	0.147
White Sucker	1	1.587	0.051	Mottled Sculpin	12	11.650	0.879
				Rainbow Trout	9	8.738	0.659
Spring 2005	7/5/2005			Summer 2005	8/9/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	24	43.636	1.341	Blacknose Dace	52	61.905	5.652
Brook Stickleback	1	1.818	0.056	Brook Stickleback	2	2.381	0.217
Brook Trout	1	1.818	0.056	Brook Trout	1	1.190	0.109
Brown Trout	2	3.636	0.112	Brown Trout	3	3.571	0.326
Burbot	1	1.818	0.056	Central Mudminnow	6	7.143	0.652
Central Mudminnow	8	14.545	0.447	Coho Salmon	2	2.381	0.217
Creek Chub	3	5.455	0.168	Creek Chub	6	7.143	0.652
Mottled Sculpin	10	18.182	0.559	Johnny Darter	2	2.381	0.217
Northern Redbelly Dace	2	3.636	0.112	Mottled Sculpin	1	1.190	0.109
Rainbow Trout	2	3.636	0.112	Rainbow Trout	7	8.333	0.761
				White Sucker	2	2.381	0.217

Steinberg Road Dow	nstream #2						
Spring 2004	7/16/2004			Summer 2004	8/16/2004		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	50	44.643	2.101	Blacknose Dace	63	56.250	4.484
Bluegill	1	0.893	0.042	Bluegill	1	0.893	0.071
Brown Trout	8	7.143	0.336	Brook Trout	1	0.893	0.071
Central Mudminnow	2	1.786	0.084	Brown Trout	5	4.464	0.356
Coho Salmon	2	1.786	0.084	Central Mudminnow	10	8.929	0.712
Creek Chub	7	6.250	0.294	Chinook Salmon	1	0.893	0.071
Longnose Dace	22	19.643	0.924	Coho Salmon	3	2.679	0.214
Mottled Sculpin	14	12.500	0.588	Creek Chub	5	4.464	0.356
Rainbow Trout	6	5.357	0.252	Mottled Sculpin	7	6.250	0.498
				Pumpkinseed	1	0.893	0.071
				Rainbow Trout	12	10.714	0.854
				White Sucker	3	2.679	0.214
Spring 2005	7/5/2005			Summer 2005	8/9/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	18		1.071	Blacknose Dace	39	42.857	4.239
Brook Stickleback	1	2.500	0.060	Brook Trout	2	2.198	0.217
Brook Trout	1	2.500	0.060	Brown Trout	5	5.495	0.543
Brown Trout	3	7.500	0.179	Central Mudminnow	5	5.495	0.543
Central Mudminnow	4	10.000	0.238	Chinook Salmon	1	1.099	0.109
Creek Chub	1	2.500	0.060	Coho Salmon	5	5.495	0.543
Mottled Sculpin	9	22.500	0.536	Creek Chub	1	1.099	0.109
Rainbow Trout	2	5.000	0.119	Mottled Sculpin	19	20.879	2.065
White Sucker	1	2.500	0.060	Rainbow Trout	12	13.187	1.304
				White Sucker	2	2.198	0.217

0 ' 000'	nstream #3						
Spring 2004	7/16/2004			Summer 2004	8/16/2004		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	50	43.103	1.609	Blacknose Dace	61	48.8	4.480
Bluegill	1	0.862	0.032	Bluegill	1	0.8	0.073
Brown Trout	18	15.517	0.579	Brook Trout	1	0.8	0.073
Central Mudminnow	1	0.862	0.032	Brown Trout	23	18.4	1.689
Creek Chub	4	3.448	0.129	Chinook Salmon	1	0.8	0.073
Longnose Dace	2	1.724	0.064	Coho Salmon	4	3.2	0.294
Mottled Sculpin	31	26.724	0.998	Creek Chub	6	4.8	0.441
Rainbow Trout	9	7.759	0.290	Mottled Sculpin	17	13.6	1.248
				Rainbow Trout	11	8.8	0.808
Spring 2005	7/5/2005	;		Summer 2005	8/9/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	19	30.645	1.132	Blacknose Dace	59	32.240	4.487
Brown Trout	2	3.226	0.119	Brown Trout	7	3.825	0.532
Central Mudminnow	2	3.226	0.119	Chinook Salmon	5	2.732	0.380
Coho Salmon	1	1.613	0.060	Coho Salmon	3	1.639	0.228
Mottled Sculpin	32	51.613	1.907	Creek Chub	1	0.546	0.076
Northern Redbelly Da	ice 1	1.613	0.060	Johnny Darter	2	1.093	0.152
Rainbow Trout	5	8.065	0.298	Mottled Sculpin	85	46.448	6.464
				Rainbow Trout	17	9.290	1.293
				White Sucker	4	2.186	0.304
Huff Road							
Tiuli Road							
Spring 2004	5/19/2004			Summer 2004	8/17/2004		
Spring 2004	5/19/2004 Number P	roportion Ca	atch/Minute	Summer 2004	8/17/2004 Number		Catch/Minute
Species	Number P	roportion Ca		Species	Number	Proportion	Catch/Minute
Species Bluegill	Number P	0.844	0.024	Species Bluegill	Number 3	Proportion 0.513	0.041
Species Bluegill Brown Trout	Number P 2 93	0.844 39.241	0.024 1.130	Species Bluegill Brown Trout	Number 3 292	Proportion 0.513 49.915	0.041 4.031
Species Bluegill Brown Trout Central Mudminnow	Number P 2 93 1	0.844 39.241 0.422	0.024 1.130 0.012	Species Bluegill Brown Trout Chinook Salmon	Number 3 292 6	Proportion 0.513 49.915 1.026	0.041 4.031 0.083
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon	Number P 2 93 1 25	0.844 39.241 0.422 10.549	0.024 1.130 0.012 0.304	Species Bluegill Brown Trout Chinook Salmon Coho Salmon	Number 3 292 6 19	Proportion 0.513 49.915 1.026 3.248	0.041 4.031 0.083 0.262
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon	Number P 2 93 1 25 23	0.844 39.241 0.422 10.549 9.705	0.024 1.130 0.012 0.304 0.280	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin	Number 3 292 6 19 68	Proportion 0.513 49.915 1.026 3.248 11.624	0.041 4.031 0.083 0.262 0.939
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner	Number P 2 93 1 25 23 2	0.844 39.241 0.422 10.549 9.705 0.844	0.024 1.130 0.012 0.304 0.280 0.024	Species Bluegill Brown Trout Chinook Salmon Coho Salmon	Number 3 292 6 19	Proportion 0.513 49.915 1.026 3.248 11.624	0.041 4.031 0.083 0.262 0.939
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin	Number P 2 93 1 25 23 2 46	0.844 39.241 0.422 10.549 9.705 0.844 19.409	0.024 1.130 0.012 0.304 0.280 0.024 0.559	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin	Number 3 292 6 19 68	Proportion 0.513 49.915 1.026 3.248 11.624	0.041 4.031 0.083 0.262 0.939
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner	Number P 2 93 1 25 23 2	0.844 39.241 0.422 10.549 9.705 0.844	0.024 1.130 0.012 0.304 0.280 0.024	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin	Number 3 292 6 19 68	Proportion 0.513 49.915 1.026 3.248 11.624	0.041 4.031 0.083 0.262 0.939
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead	Number P 2 93 1 25 23 2 46 44 1	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout	Number 3 292 6 19 68 197	Proportion 0.513 49.915 1.026 3.248 11.624 33.675	0.041 4.031 0.083 0.262 0.939
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead Spring 2005	Number P 2 93 1 25 23 2 46 44 1 6/10/2005	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565 0.422	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535 0.012	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout	Number 3 292 6 19 68 197	Proportion 0.513 49.915 1.026 3.248 11.624 33.675	0.041 4.031 0.083 0.262 0.939 2.720
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead Spring 2005 Species	Number P 2 93 1 25 23 2 46 44 1 6/10/2005 Number P	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565 0.422	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535 0.012	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout Summer 2005 Species	Number 3 292 6 19 68 197 8/8/2005 Number	Proportion 0.513 49.915 1.026 3.248 11.624 33.675	0.041 4.031 0.083 0.262 0.939 2.720
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead Spring 2005 Species Brook Trout	Number P 2 93 1 25 23 2 46 44 1 6/10/2005 Number P	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565 0.422	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535 0.012	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout Summer 2005 Species Brown Trout	Number 3 292 6 19 68 197 8/8/2005 Number 244	Proportion 0.513 49.915 1.026 3.248 11.624 33.675 Proportion 50.206	0.041 4.031 0.083 0.262 0.939 2.720 Catch/Minute 5.014
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead Spring 2005 Species Brook Trout Brown Trout	Number P 2 93 1 25 23 2 46 44 1 6/10/2005 Number P 3 135	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565 0.422 roportion Ca 1.075 48.387	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535 0.012 atch/Minute 0.051 2.274	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout Summer 2005 Species Brown Trout Burbot	Number 3 292 6 19 68 197 8/8/2005 Number 244 3	Proportion 0.513 49.915 1.026 3.248 11.624 33.675 Proportion 50.206 0.617	0.041 4.031 0.083 0.262 0.939 2.720 Catch/Minute 5.014 0.062
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead Spring 2005 Species Brook Trout Brown Trout Chinook Salmon	Number P 2 93 1 25 23 2 46 44 1 6/10/2005 Number P 3 135 21	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565 0.422 roportion Ca 1.075 48.387 7.527	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535 0.012 atch/Minute 0.051 2.274 0.354	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout Summer 2005 Species Brown Trout Burbot Chinook Salmon	Number 3 292 6 19 68 197 8/8/2005 Number 244 3 14	Proportion 0.513 49.915 1.026 3.248 11.624 33.675 Proportion 50.206 0.617 2.881	0.041 4.031 0.083 0.262 0.939 2.720 Catch/Minute 5.014 0.062 0.288
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead Spring 2005 Species Brook Trout Brown Trout Chinook Salmon Coho Salmon	Number P 2 93 1 25 23 2 46 44 1 6/10/2005 Number P 3 135 21 5	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565 0.422 roportion Ca 1.075 48.387 7.527 1.792	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535 0.012 atch/Minute 0.051 2.274 0.354 0.084	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout Summer 2005 Species Brown Trout Burbot Chinook Salmon Coho Salmon	Number 3 292 6 19 68 197 8/8/2005 Number 244 3 14	Proportion 0.513 49.915 1.026 3.248 11.624 33.675 Proportion 50.206 0.617 2.881 3.292	0.041 4.031 0.083 0.262 0.939 2.720 Catch/Minute 5.014 0.062 0.288 0.329
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead Spring 2005 Species Brook Trout Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin	Number P 2 93 1 25 23 2 46 44 1 6/10/2005 Number P 3 135 21 5 40	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565 0.422 roportion Ca 1.075 48.387 7.527 1.792 14.337	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535 0.012 atch/Minute 0.051 2.274 0.354 0.084 0.674	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout Summer 2005 Species Brown Trout Burbot Chinook Salmon Coho Salmon Creek Chub	Number 3 292 6 19 68 197 8/8/2005 Number 244 3 14 16 1	Proportion 0.513 49.915 1.026 3.248 11.624 33.675 Proportion 50.206 0.617 2.881 3.292 0.206	0.041 4.031 0.083 0.262 0.939 2.720 Catch/Minute 5.014 0.062 0.288 0.329 0.021
Species Bluegill Brown Trout Central Mudminnow Chinook Salmon Coho Salmon Common Shiner Mottled Sculpin Rainbow Trout Yellow Bullhead Spring 2005 Species Brook Trout Brown Trout Chinook Salmon Coho Salmon	Number P 2 93 1 25 23 2 46 44 1 6/10/2005 Number P 3 135 21 5	0.844 39.241 0.422 10.549 9.705 0.844 19.409 18.565 0.422 roportion Ca 1.075 48.387 7.527 1.792	0.024 1.130 0.012 0.304 0.280 0.024 0.559 0.535 0.012 atch/Minute 0.051 2.274 0.354 0.084	Species Bluegill Brown Trout Chinook Salmon Coho Salmon Mottled Sculpin Rainbow Trout Summer 2005 Species Brown Trout Burbot Chinook Salmon Coho Salmon	Number 3 292 6 19 68 197 8/8/2005 Number 244 3 14	Proportion 0.513 49.915 1.026 3.248 11.624 33.675 Proportion 50.206 0.617 2.881 3.292 0.206 11.111	0.041 4.031 0.083 0.262 0.939 2.720 Catch/Minute 5.014 0.062 0.288 0.329 0.021 1.110

Bear Creek Electrofishing Data

Milks Road Upstream							
Summer 2004	8/8/2004			Spring 2005	6/22/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minu
Blacknose Dace	105	41.339	1.881	Blacknose Dace	139	41.742	2.089
Bluegill	7	2.756	0.125	Bluegill	1	0.300	0.015
Brook Stickleback	1	0.394	0.018	Brook Trout	7	2.102	0.105
Brown Trout	11	4.331	0.197	Brown Trout	8	2.402	0.120
Burbot	11	4.331	0.197	Burbot	3	0.901	0.045
Central Mudminnow	1	0.394	0.018	Central Mudminnow	9	2.703	0.135
Chinook Salmon	1	0.394	0.018	Chinook Salmon	44	13.213	0.661
Creek Chub	16	6.299	0.287	Coho Salmon	4	1.201	0.060
Johnny Darter	10	3.937	0.179	Creek Chub	1	0.300	0.015
Longnose Dace	10	3.937	0.179	Golden Redhorse	1	0.300	0.015
Mottled Sculpin	12	4.724	0.215	Johnny Darter	13	3.904	0.195
Northern Pike	1	0.394	0.018	Lamprey	2	0.601	0.030
Rainbow Trout	19	7.480	0.340	Mottled Sculpin	12	3.604	0.180
White Sucker	49	19.291	0.878	Rainbow Trout	69	20.721	1.037
				White Sucker	20	6.006	0.301
Summer 2005	8/3/2005						
Species			Catch/Minute				
Blacknose Dace	156	32.099	2.818				
Brook Trout	4	0.823	0.072				
Brown Trout	14	2.881	0.253				
Burbot	4	0.823	0.072				
Central Mudminnow	7	1.440	0.126				
Chinook Salmon	9	1.852	0.163				
Coho Salmon	3	0.617	0.054				
Creek Chub	14	2.881	0.253				
Green Sunfish	1	0.206	0.018				
Johnny Darter	37	7.613	0.668				
Mottled Sculpin	62	12.757	1.120				
Northern Brook Lamprey	7	1.440	0.126				
Rainbow Trout	127	26.132	2.294				
White Sucker	40	8.230	0.723				
Yellow Perch	1	0.206	0.018				

Milks Road Downstream							
Summer 2004	8/8/2004			Spring 2005	6/22/2005		
Species	Number F	Proportion (Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	137	49.104	2.394	Blacknose Dace	88	33.208	1.597
Bluegill	27	9.677	0.472	Brook Trout	1	0.377	0.018
Brown Trout	8	2.867	0.140	Brown Bullhead	1	0.377	0.018
Burbot	2	0.717	0.035	Brown Trout	11	4.151	0.200
Central Mudminnow	2	0.717	0.035	Burbot	2	0.755	0.036
Chinook Salmon	2	0.717	0.035	Central Mudminnow	2	0.755	0.036
Creek Chub	29	10.394	0.507	Chinook Salmon	53	20.000	0.962
Green Sunfish	2	0.717	0.035	Coho Salmon	2	0.755	0.036
Johnny Darter	12	4.301	0.210	Creek Chub	11	4.151	0.200
Mottled Sculpin	25	8.961	0.437	Golden Redhorse	1	0.377	0.018
Rainbow Trout	17	6.093	0.297	Johnny Darter	6	2.264	0.109
White Sucker	16	5.735	0.280	Lamprey	3	1.132	0.054
				Mottled Sculpin	8	3.019	0.145
				Rainbow Trout	62	23.396	1.125
				White Sucker	14	5.283	0.254
Summer 2005	8/3/2005						
Species	Number F	Proportion (Catch/Minute				
Blacknose Dace	67	25.475	1.475				
Bluntnose Minnow	2	0.760	0.044				
Brook Trout	1	0.380	0.022				
Brown Trout	6	2.281	0.132				
Burbot	13	4.943	0.286				
Central Mudminnow	3	1.141	0.066				
Chinook Salmon	11	4.183	0.242				
Coho Salmon	9	3.422	0.198				
Creek Chub	12	4.563	0.264				
Green Sunfish	2	0.760	0.044				
Johnny Darter	12	4.563	0.264				
Largemouth Bass	1	0.380	0.022				
Longnose Dace	2	0.760	0.044				
Mottled Sculpin	24	9.125	0.528				
Northern Brook Lamprey	5	1.901	0.110				
Pumpkinseed	1	0.380	0.022				
Rainbow Trout	77	29.278	1.695				
White Sucker	14	5.323	0.308				
Yellow Perch	1	0.380	0.022				

Summer 2004	8/6/2004			Spring 2005	6/7/2005		
Species	Number	Proportion	Catch/Minute	Species	Number	Proportion	Catch/Minute
Blacknose Dace	139	50.91575	3.682119205	Blacknose Dace	73	32.444	2.02
Bluegill	6	2.197802	0.158940397	Bluegill	1	0.444	0.02
Brown Trout	13	4.761905	0.344370861	Brook Trout	2	0.889	0.05
Burbot	2	0.732601	0.052980132	Brown Trout	1	0.444	0.02
Central Mudminnow	1	0.3663	0.026490066	Burbot	5	2.222	0.13
Chinook Salmon	2	0.732601	0.052980132	Chinook Salmon	72	32.000	1.99
Creek Chub	17	6.227106	0.450331126	Coho Salmon	6	2.667	0.16
Golden Redhorse Sucker	1	0.3663	0.026490066	Creek Chub	9	4.000	0.24
Johnny Darter	19	6.959707	0.503311258	Golden Redhorse	1	0.444	0.02
Longnose Dace	3	1.098901	0.079470199	Johnny Darter	8	3.556	0.22
Mottled Sculpin	7	2.564103	0.185430464	Mottled Sculpin	7	3.111	0.19
Rainbow Trout	32	11.72161	0.847682119	Rainbow Trout	22	9.778	0.60
Shorthead Redhorse Sucker	3	1.098901	0.079470199	White Sucker	18	8.000	0.49
White Sucker	28	10.25641	0.741721854				
Summer 2005	8/2/2005						
Species	Number	Proportion	Catch/Minute				
Blacknose Dace	85	30.576	2.044				
Bluegill	1	0.360	0.024				
Bluntnose Minnow	1	0.360	0.024				
Brown Trout	9	3.237	0.216				
Burbot	8	2.878	0.192				
Chinook Salmon	13	4.676	0.313				
Coho Salmon	6	2.158	0.144				
Creek Chub	9	3.237	0.216				
Golden Redhorse Sucker	2	0.719	0.048				
Johnny Darter	10	3.597	0.240				
Largemouth Bass	1	0.360	0.024				
Longnose Dace	1	0.360	0.024				
Mottled Sculpin	26	9.353	0.625				
Northern Brook Lamprey	4	1.439	0.096				
Northern Redbelly Dace	4	1.439	0.096				
Rainbow Trout	87	31.295	2.092				
White Sucker	11	3.957	0.265				

Swain Property Downstream							
Summer 2004	8/6/2004			Spring 2005	6/7/2005		
Species			Catch/Minute	Species	Number		Catch/Minute
Blacknose Dace	165	49.107		Blacknose Dace	26	22.609	0.783
Bluegill	1	0.298	0.026	Brook Trout	1	0.870	0.030
Brook Stickleback	1	0.298	0.026	Brown Trout	1	0.870	0.030
Brown Trout	5	1.488	0.132	Central Mudminnow	2	1.739	0.060
Burbot	1	0.298	0.026	Chinook Salmon	30	26.087	0.904
Central Mudminnow	5	1.488	0.132	Coho Salmon	3	2.609	0.090
Chinook Salmon	2	0.595	0.053	Creek Chub	4	3.478	0.120
Coho Salmon	5	1.488	0.132	Golden Redhorse	12	10.435	0.361
Common Shiner	3	0.893	0.079	Johnny Darter	11	9.565	0.331
Creek Chub	19	5.655	0.503	Mottled Sculpin	1	0.870	0.030
Golden Redhorse Sucker	1	0.298	0.026	Rainbow Trout	8	6.957	0.241
Johnny Darter	22	6.548	0.583	White Sucker	16	13.913	0.482
Longnose Dace	1	0.298	0.026				
Mottled Sculpin	15	4.464	0.397				
Rainbow Trout	65	19.345	1.722				
Shorthead Redhorse Sucker	1	0.298	0.026				
White Sucker	24	7.143	0.636				
Summer 2005	8/2/2005						
Species	Number	Proportion	Catch/Minute				
Blacknose Dace	59	32.597	1.674				
Brown Trout	9	4.972	0.255				
Burbot	3	1.657	0.085				
Chinook Salmon	7	3.867	0.199				
Creek Chub	2	1.105	0.057				
Golden Redhorse Sucker	3	1.657	0.085				
Johnny Darter	14	7.735	0.397				
Largemouth Bass	3	1.657	0.085				
Mottled Sculpin	10	5.525	0.284				
Northern Brook Lamprey	2	1.105	0.057				
Rainbow Trout	60	33.149					
White Sucker	9	4.972	0.255				

Johnson Road							
Summer 2004	7/30/2004			Spring 2005	6/3/2005		
Species			Catch/Minute	Species	Number	Proportion C	
Blacknose Dace	237	50.641	2.841	Blacknose Dace	66		1.018
Bluegill	7	1.496	0.084	Bluegill	2		0.031
Brook Stickleback	5	1.068	0.060	Brown Trout	5		0.077
Brown Trout	6	1.282	0.072	Central Mudminnow	1	0.410	0.015
Burbot	1	0.214	0.012	Chinook Salmon	22		0.339
Central Mudminnow	18	3.846	0.216	Coho Salmon	27	11.066	0.417
Coho Salmon	5	1.068	0.060	Common Shiner	3		0.046
Common Shiner	1	0.214	0.012	Creek Chub	13	5.328	0.201
Creek Chub	25	5.342	0.300	Golden Redhorse	5	2.049	0.077
Greater Redhorse Sucker	1	0.214	0.012	Johnny Darter	15	6.148	0.231
Johnny Darter	15	3.205	0.180	Longnose Dace	1	0.410	0.015
Longnose Dace	29	6.197	0.348	Mottled Sculpin	11	4.508	0.170
Mottled Sculpin	41	8.761	0.491	Northern Redbelly Dace	4	1.639	0.062
Raibow Trout	37	7.906	0.443	Rainbow Trout	15	6.148	0.231
White Sucker	40	8.547	0.479	White Sucker	54	22.131	0.833
Summer 2005	8/5/2005						
Species	Number F	Proportion C	Catch/Minute				
Blacknose Dace	225	27.881	3.238				
Bluegill	1	0.124	0.014				
Brown Trout	9	1.115	0.130				
Burbot	3	0.372	0.043				
Central Mudminnow	9	1.115	0.130				
Chinook Salmon	95	11.772	1.367				
Common Shiner	4	0.496	0.058				
Creek Chub	45	5.576	0.648				
Golden Redhorse Sucker	1	0.124	0.014				
Johnny Darter	39	4.833	0.561				
Largemouth Bass	3	0.372	0.043				
Longnose Dace	13	1.611	0.187				
Mottled Sculpin	69	8.550	0.993				
Northern Redbelly Dace	5	0.620	0.072				
Rainbow Trout	227	28.129	3.267				
Redear Sunfish	1	0.124	0.014				
White Sucker	58	7.187	0.835				

Spirit of the Woods							
Summer 2004	7/28/2004			Spring 2005	6/2/2005		
Species	Number		Catch/Minute	Species	Number	•	Catch/Minu
Blacknose Dace	70			Blacknose Dace	10		0.165
Bluegill	8	1.848	0.077	Brown Trout	26	14.054	0.430
Brown Trout	48		0.460	Burbot	11		0.182
Burbot	72		0.690	Chinook Salmon	15		0.248
Central Mudminnow	3	0.693	0.029	Coho Salmon	5	2.703	0.083
Coho Salmon	4	0.924	0.038	Creek Chub	6	3.243	0.099
Common Shiner	1	0.231	0.010	Golden Redhorse	5	2.703	0.083
Creek Chub	99	22.864	0.949	Johnny Darter	6	3.243	0.099
Greater Redhorse Sucker	2	0.462	0.019	Longnose Dace	1	0.541	0.017
Johnny Darter	7	1.617	0.067	Mottled Sculpin	5	2.703	0.083
Longnose Dace	27	6.236	0.259	Pumpkinseed	1	0.541	0.017
Mottled Sculpin	25	5.774	0.240	Rainbow Trout	14	7.568	0.231
Rainbow Trout	21	4.850	0.201	Shorthead Redhorse	23	12.432	0.380
White Sucker	45	10.393	0.431	White Sucker	56	30.270	0.925
Yellow Perch	1	0.231	0.010	Yellow Perch	1	0.541	0.017
Summer 2005	8/4/2005						
Species	Number	Proportion	Catch/Minute				
Blacknose Dace	11	7.534	0.181				
Brown Trout	30	20.548	0.493				
Chinook Salmon	14	9.589	0.230				
Coho Salmon	3	2.055	0.049				
Common Shiner	2	1.370	0.033				
Creek Chub	11	7.534	0.181				
Golden Redhorse Sucker	2	1.370	0.033				
Johnny Darter	2	1.370	0.033				
Mottled Sculpin	3	2.055	0.049				
Rainbow Trout	22	15.068	0.362				
Redear Sunfish	1	0.685	0.016				
White Sucker	45	30.822	0.740				

Summer 2004	7/27/2004			Spring 2005	6/1/2005		
Species	Number	Proportion C		Species	Number		Catch/Minute
Alewife	2	0.548	0.025	Blacknose Dace	10		0.159
Blacknose Dace	1	0.274	0.013	Brown Trout	14		0.223
Bluegill	1	0.274	0.013	Burbot	47	15.878	0.748
Brown Trout	16	4.384	0.202	Central Mudminnow	29	9.797	0.462
Burbot	217	59.452	2.738	Chinook Salmon	30	10.135	0.477
Central Mudminnow	21	5.753	0.265	Coho Salmon	21	7.095	0.334
Chinook Salmon	2	0.548	0.025	Common Shiner	2	0.676	0.032
Coho Salmon	1	0.274	0.013	Creek Chub	9	3.041	0.143
Creek Chub	12	3.288	0.151	Johnny Darter	3	1.014	0.048
Grass Pickerel	1	0.274	0.013	Mottled Sculpin	39	13.176	0.621
Johnny Darter	25	6.849	0.315	Rainbow Trout	20	6.757	0.318
Mottled Sculpin	37	10.137	0.467	Shorthead Redhorse	1	0.338	0.016
Northern Pike	1	0.274	0.013	White Sucker	71	23.986	1.130
Rainbow Trout	5	1.370	0.063				
Rock Bass	7	1.918	0.088				
White Sucker	16	4.384	0.202				
Summer 2005	8/1/2005						
Species	Number	Proportion C	Catch/Minute				
Blacknose Dace	3	1.079	0.059				
Blacksided Dace	2	0.719	0.039				
Brown Trout	13	4.676	0.254				
Burbot	103	37.050	2.014				
Central Mudminnow	27	9.712	0.528				
Chinook Salmon	12	4.317	0.235				
Common Shiner	1	0.360	0.020				
Creek Chub	32	11.511	0.626				
Johnny Darter	10	3.597	0.196				
Largemouth Bass	1	0.360	0.020				
Mottled Sculpin	24	8.633	0.469				
Northern Brook Lamprey	1	0.360	0.020				
Northern Pike	2	0.719	0.039				
Rainbow Trout	14	5.036	0.274				
Rock Bass	6	2.158	0.117				
White Sucker	27	9.712	0.528				