

4-2008

## **A Carboxyl-terminal Sequence in the Lutropin $\beta$ Subunit Contributes to the Sorting of Lutropin to the Regulated Pathway**

Albina Jablonka-Shariff  
*Washington University School of Medicine in St. Louis*

Christopher A. Pearl  
*Grand Valley State University, pearlch@gvsu.edu*

Anna Comstock  
*Washington University School of Medicine in St. Louis*

Irving Boime  
*Washington University School of Medicine in St. Louis*

Follow this and additional works at: [https://scholarworks.gvsu.edu/bms\\_articles](https://scholarworks.gvsu.edu/bms_articles)



Part of the [Endocrinology Commons](#)

---

### **ScholarWorks Citation**

Jablonka-Shariff, Albina; Pearl, Christopher A.; Comstock, Anna; and Boime, Irving, "A Carboxyl-terminal Sequence in the Lutropin  $\beta$  Subunit Contributes to the Sorting of Lutropin to the Regulated Pathway" (2008). *Peer Reviewed Articles*. 62.

[https://scholarworks.gvsu.edu/bms\\_articles/62](https://scholarworks.gvsu.edu/bms_articles/62)

This Article is brought to you for free and open access by the Biomedical Sciences Department at ScholarWorks@GVSU. It has been accepted for inclusion in Peer Reviewed Articles by an authorized administrator of ScholarWorks@GVSU. For more information, please contact [scholarworks@gvsu.edu](mailto:scholarworks@gvsu.edu).

# A Carboxyl-terminal Sequence in the Lutropin $\beta$ Subunit Contributes to the Sorting of Lutropin to the Regulated Pathway\*

Received for publication, January 24, 2008, and in revised form, February 20, 2008. Published, JBC Papers in Press, February 21, 2008, DOI 10.1074/jbc.M800654200

Albina Jablonka-Shariff<sup>1</sup>, Christopher A. Pearl<sup>1,2</sup>, Anna Comstock, and Irving Boime<sup>3</sup>

From the Department of Molecular Biology and Pharmacology, Washington University School of Medicine, St. Louis, Missouri 63110

Although synthesized in the same pituitary gonadotropes, the secretion profiles of lutropin (LH) and follitropin (FSH) differ. LH is secreted through a regulated pathway and associated with a bolus release at mid-estrous cycle. In contrast, the majority of FSH is secreted constitutively with an incremental increase until ovulation. Both share an identical  $\alpha$  subunit, and thus the  $\beta$  subunit contains determinants for sorting into the regulated pathway. Previously, we demonstrated that a hydrophobic carboxyl-terminal heptapeptide of the LH $\beta$  subunit (Leu-Ser-Gly-Leu-Leu-Phe-Leu), not found in the FSH $\beta$  subunit, influences the intracellular behavior of the LH dimer. To test the hypothesis that the peptide contributes to differential sorting, we monitored the fates of LH and LH $\Delta$ T (LH $\beta$  subunit lacking the carboxyl-terminal seven amino acids) dimers in the rat somatotrope-derived GH<sub>3</sub> cell line in which both the regulated and constitutive secretory pathways operate. Pulse-chase labeling demonstrated that the LH $\Delta$ T dimer was diverted to the constitutive pathway, resulting in a significant decrease in the corresponding intracellular pool. Forskolin stimulated LH dimer release 3-fold, which was accompanied by a parallel decrease of intracellular LH; only marginal forskolin stimulation of LH $\Delta$ T was seen. Immunofluorescence after cycloheximide treatment demonstrated decreased retention of LH $\Delta$ T compared with LH, consistent with increased constitutive secretion of LH $\Delta$ T. We also demonstrated that fusing the heptapeptide to the carboxyl terminus of the FSH $\beta$  subunit resulted in an increased regulated secretion of this FSH analog compared with wild-type FSH. These data are the first to identify a novel structural determinant responsible for the sorting of a member of the glycoprotein hormone family into the regulated secretory pathway.

Lutropin (LH)<sup>4</sup> and follitropin (FSH) are synthesized and secreted by pituitary gonadotropes and are members of the gly-

coprotein hormone family, which also includes thyrotropin (TSH) and the placental hormone chorionic gonadotropin (CG). They are heterodimers that share a common  $\alpha$  subunit but differ in their hormone-specific  $\beta$  subunits (1, 2). Both subunits are glycosylated, containing asparagine (N)-linked oligosaccharides (3, 4). The mature carbohydrate structures are hormone-specific in that the terminal oligosaccharide is sulfate for LH (3, 5), whereas FSH contains sialic acid (3).

LH and FSH play key roles in regulating reproductive function. In females, FSH stimulates follicular growth, maintaining a steady concentration during the early follicular stage, and is required to facilitate selection of follicles to the preovulatory phase. At this time, low levels of LH stimulate steroidogenesis in thecal cells by enhancing androgen synthesis, which in turn is converted to estradiol in the presence of FSH. The gradual increase in estradiol is essential to initiate the LH surge. The reciprocal relationship between FSH and estrogen concentrations during the follicular phase of the menstrual cycle is an exquisitely sensitive feedback pathway that governs the selection of the preovulatory follicle (6, 7). The modes of secretion for LH and FSH are linked to their function; LH is released in pulses via a regulated pathway (*i.e.* LH is stored in secretory granules) and associated with a bolus release at midcycle to rupture the follicle and form the corpus luteum (6–9). By contrast, FSH is primarily constitutively secreted, tightly coupled to its synthesis rate with gradual incremental increases until ovulation (9, 10).

The N-linked oligosaccharides play a critical role in the extracellular stability of both hormones. Sulfated oligosaccharides lead to a rapid clearance of LH *in vivo*, regulating its pulsatile release (3, 10, 11). Sialylation results in greater extracellular longevity of FSH as compared with LH. Although the structural motifs that govern the differential sorting of LH and FSH have yet to be identified, critical structural cues exist that might account for these intracellular events. LH and FSH are synthesized in the same cell and share an identical  $\alpha$  subunit, and thus the  $\beta$  subunit must represent the key determinant for the specificity of carbohydrate processing and for the intracellular segregation of one or both hormones. Of all the human glycoprotein hormone  $\beta$  subunits, the LH $\beta$  subunit is the most hydrophobic, particularly in the region between residues 75 and 121 (12, 13). DNA sequences for the LH $\beta$  (12) and thyrotropin  $\beta$  (14) subunits encode hydrophobic stretches of seven and six amino acids, respectively, at their carboxyl termini, but a similar heptapeptide is not observed at the carboxyl terminus of the FSH $\beta$  subunit. This difference points to the carboxyl end

\* This manuscript was supported in part by National Institutes of Health (NIH) Grant DR065155, NIH Neuroscience Blueprint Core Grant NS057105 to Washington University, and the Bakewell Family Foundation. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> Both authors contributed equally to this manuscript.

<sup>2</sup> Supported by Training Grant T32HD049305-03 from the National Institutes of Health/NICHD.

<sup>3</sup> To whom correspondence should be addressed: 660 S. Euclid Ave., Box 8103, St. Louis, MO 63110. Tel.: 314-362-2556; Fax: 314-362-7051; E-mail: iboime@wustl.edu.

<sup>4</sup> The abbreviations used are: LH, lutropin (luteinizing hormone); FSH, follitropin (follicle-stimulating hormone); CG, chorionic gonadotropin; CHX, cycloheximide; ER, endoplasmic reticulum; CHO, Chinese hamster ovary.

## Sorting Signal for LH

of the LH $\beta$  subunit as a potential candidate for a sorting determinant.

Some important observations that might explain the unique secretion patterns discussed above have been obtained from transfected animal cell lines. Earlier studies from our laboratory and others demonstrated that the unassembled pituitary  $\beta$  subunits do not efficiently exit the ER in the absence of the  $\alpha$  subunit (15–17). Although co-expression with the  $\alpha$  subunit rescues the  $\beta$  subunits, differences exist in the extent of assembly of the  $\alpha/\beta$  subunit pairs. For example, in the case of LH $\beta$ , the amount of dimer formed in transfected CHO cells is less than 10% (12, 15, 18), whereas more than 80% of the steady-state FSH $\beta$  subunit is secreted as a component of the heterodimer (16). The terminal LH $\beta$  heptapeptide (Leu-Ser-Gly-Leu-Leu-Phe-Leu) accounts, in part, for this inefficient assembly (13, 18). Based on these observations, we proposed that this sequence serves as a signal capable of governing the intracellular sorting and trafficking of LH (13).

The experiments described above were performed with CHO cells, which secrete proteins only by the constitutive route (13, 16–19) without intracellular accumulation of mature hormone dimers, thus precluding studies of glycoprotein hormone secretion by the regulated pathway. To investigate sorting of the LH/FSH dimers and free subunits, we have used the GH<sub>3</sub> cell line (20, 21), which is derived from pituitary somatotropes and contain storage vesicles responsive to secretagogues (20–22). Importantly, we previously demonstrated that transfected GH<sub>3</sub> cells secrete LH and FSH primarily through regulated and constitutive secretory pathways, respectively (21), and that the LH *N*-linked carbohydrates were sulfated (20). The above observations were in marked contrast with data obtained from transfected CHO cells in which no detectable responses to secretagogues or LH sulfation were observed (20, 21). Here we tested the requirement of the LH $\beta$  heptapeptide to direct LH to the regulated pathway of GH<sub>3</sub> cells. The data indicate that the carboxyl-terminal heptapeptide contributes to the trafficking of LH dimer to the regulated secretory pathway.

## EXPERIMENTAL PROCEDURES

**Cell Culture and Stable Transfection**—GH<sub>3</sub> cells were a gift from Dr. Dennis Shields (Albert Einstein College of Medicine, New York). The cells were grown (no more than 30 passages) at 37 °C in Ham's F-12 medium (Mediatech Inc., Herndon, VA) supplemented with 12.5% horse serum (Invitrogen), 2.5% fetal bovine serum (Harlan Bioproducts for Science, Inc., Indianapolis, IN), 2 mM L-glutamine, 100 units/ml penicillin, and 100  $\mu$ g/ml streptomycin in a humidified 5% CO<sub>2</sub> incubator. Transfections were performed using Lipofectamine 2000 (Invitrogen) on semiconfluent cells in 6-well plates. Cells were transfected with 4  $\mu$ g of the  $\alpha$ , LH $\beta$ , LH $\beta$ 114 (designated LH $\beta$  $\Delta$ T), FSH $\beta$ , or FSH $\beta$ -LH $\beta$  chimera (designated FSH $\beta$ -L) subunit genes (Fig. 1) contained in the vector pM<sup>2</sup>HA (19) to obtain clones expressing the individual subunits and corresponding dimers. Stable clones were selected ~16 days later with 0.25 mg/ml G418 (Research Product International, Mt. Prospect, IL). Single colonies were isolated and subsequently screened by immunoprecipitating the media and lysates of metabolically labeled cells (see below). Several clones ( $n = 5$  per dimer)

expressing the dimers LH, LH $\Delta$ T, FSH, or FSH-L were maintained in culture in the presence of 0.125 mg/ml G418 and used for the experiments described below.

The mutant LH $\beta$  $\Delta$ T described previously lacks a seven-amino acid extension (Leu-Ser-Gly-Leu-Leu-Phe-Leu) at the carboxyl terminus of the LH $\beta$  subunit (13). To construct FSH $\beta$ -L, the heptapeptide sequence of LH $\beta$  subunit (plus the stop codon) was inserted in-frame at the 3'-end of the FSH $\beta$  subunit by using overlapping PCR mutagenesis. The PCR was performed using KlenTag DNA polymerase (Sigma) and GenAmp PCR system 2400 (PerkinElmer Life Sciences). The following primers were used in the construction of the FSH $\beta$ -L chimera: oligo 1 (universal primer for pM<sup>2</sup>HA); 5'-TTC TCC CCC GCA GCC CTA GAA GAC GTT CCA-3'; oligo 2; 5'-GAG GAG GCC TGA GAG TTC TTT CAT TTC ACC-3'; oligo 3; 5'-GGT GAA ATG AAA GAA CTC TCA GGC CTC CTC-3'; oligo 4 (universal primer for pM<sup>2</sup>HA); 5'-TTT TCA CTG CAT TCT AGT TGT GGT TTG TCC-3'.

The universal primers (oligos 1 and 4) corresponded to the sequences in pM<sup>2</sup>HA vector located upstream and downstream of multiple cloning sites. In the first PCR reaction, the FSH $\beta$ -CTP- $\alpha$  gene was used as a template with primers 1 and 2 to amplify product A containing the entire FSH $\beta$  sequence and the beginning of the heptapeptide of LH $\beta$  subunit. A parallel reaction containing primers 3 and 4 and the LH $\beta$  subunit as a template generated PCR product B comprising part of FSH $\beta$  exon 3 and the heptapeptide sequence of LH $\beta$  with its stop codon. The overlapping PCR was performed using fragments A and B with primers 1 and 4, resulting in the final product FSH $\beta$ -L (FSH $\beta$  with the seven amino acids of LH $\beta$  subunit), which was sequenced to ensure no errors occurred during the PCR reactions. The FSH $\beta$ -L chimera was enzymatically digested by BamHI and inserted into the pM<sup>2</sup>HA vector. The pM<sup>2</sup>HA vector is a pSV2 neo derivative, which contains the ampicillin resistance and neomycin resistance genes and the Harvey murine sarcoma virus long terminal repeat.

**Metabolic Labeling and Immunoprecipitation**—For continuous labeling experiments, cells were plated into 6- or 12-well dishes and grown for ~4 days to near confluency. Cells were labeled for 16 h with 20  $\mu$ Ci/ml [<sup>35</sup>S]cysteine (specific activity > 1000 Ci/mmol, MP Biomedicals Inc., Irvine, CA) in Ham's F-12 medium minus cysteine and G418 but supplemented with 7.5% dialyzed fetal bovine serum, glutamine, and antibiotics. Labeling with inorganic sulfate was performed for 16 h in Ham's F-12 sulfate-free medium with 0.7 mCi/ml carrier-free sulfate (Na<sub>2</sub>[<sup>35</sup>S]O<sub>4</sub>; MP Biomedicals Inc.) supplemented with 7.5% dialyzed fetal bovine serum, glutamine, and antibiotics (20).

For pulse-chase experiments, confluent cells grown on 6-well plates were preincubated for 1.5 h with cysteine-free medium followed by a 20 min pulse in this medium containing 80  $\mu$ Ci/ml [<sup>35</sup>S]cysteine. At the end of the pulse, the medium was aspirated, and the cells were washed twice with prewarmed chase medium comprising Ham's F-12, 1 mM unlabeled L-cysteine (Sigma), 7.5% dialyzed fetal bovine serum, glutamine, and antibiotics and incubated in this medium for up to 24 h.

All collected media and cell lysates were treated with iodoacetamide and phenylmethanesulfonyl fluoride to inhibit pro-

teases. After centrifugation to remove cell debris, samples (2 ml) were precleared with 7.5  $\mu$ l/ml normal rabbit serum and Pansorbin (EMD Biosciences Inc., La Jolla, CA). The supernates were divided into two aliquots and immunoprecipitated for 2 h at room temperature by anti- $\alpha$  or anti-CG $\beta$  (which cross-reacts with the LH $\beta$  subunit) rabbit polyclonal sera. The immune complexes were precipitated with Pansorbin and subjected to SDS-PAGE on 12.5 or 15% gels. For quantitative comparisons of LH and LH $\Delta$ T, equal volumes of samples were always loaded on the gel. The gels were soaked in 1 M sodium salicylate for 15 min, dried, and exposed to x-ray film (18).

**Forskolin-stimulated Secretion**—To examine gonadotropin storage and regulated secretion of dimers, five wells of LH and LH $\Delta$ T cells were labeled for 16 h with [ $^{35}$ S]sulfate or [ $^{35}$ S]cysteine. Medium and lysate from one well were collected and served as the 0 time controls for the subsequent experiments. Cells in the remaining four wells were preincubated in chase medium (containing 1 mM unlabeled L-cysteine, 7.5% dialyzed fetal bovine serum, glutamine, and antibiotics) for 2 h to reduce the background of the constitutive pool. After the 2-h preincubation, media were collected and frozen for immunoprecipitation. During a second 2-h period, two wells received fresh chase medium with forskolin (25  $\mu$ M final concentration; Sigma) (20), and the remaining two wells were incubated without secretagogue. Cells expressing LH $\Delta$ T were also incubated in the presence or absence of forskolin immediately after a 16-h labeling without the 2-h preincubation period. All labeled culture media and cell lysates were then immunoprecipitated with  $\alpha$  antiserum and analyzed by 12.5% SDS-PAGE.

To examine the storage and secretion of FSH and FSH-L dimers, cells were labeled for 4 h with [ $^{35}$ S]cysteine instead of sulfate, because the N-linked oligosaccharides of FSH are not sulfated as they are in LH. Medium and lysate from one well were collected and served as the 0 time controls for the subsequent experiment. Cells in one well received chase medium for 2 h with forskolin (25  $\mu$ M final concentration) (20), and the remaining well was incubated without secretagogue. All labeled culture media and cell lysates were then immunoprecipitated with FSH dimer-specific antibody (FSH 54B) and analyzed by 15% SDS-PAGE.

**Immunofluorescence**—GH $_3$  cells expressing LH or LH $\Delta$ T were grown on Fisherbrand Superfrost-Plus microscope slides (Fisher Scientific) in Petri dishes as described above. The cells were incubated at 37  $^{\circ}$ C for 4 h in the presence or absence of cycloheximide (CHX) (15  $\mu$ g/ml; Sigma). Immediately following CHX treatment, all cells were fixed in 4% paraformaldehyde for 20 min and permeabilized with 0.2% Tween for 10 min. Cells were then incubated in 20% normal goat serum (Vector Laboratories, Burlingame, CA) for 1 h to block nonspecific binding and washed three times for 10 min in 2% bovine serum albumin (Sigma). After washing, cells were incubated in rabbit anti-CG $\beta$  antiserum (1:250; also used for immunoprecipitation experiments) for 30 min at room temperature. The cells were washed three times in 2% bovine serum albumin and incubated in goat anti-rabbit Alexa Fluor 488 (1:250; Invitrogen) for 20 min. Cells were then washed three times in 2% bovine serum albumin and once in phosphate-buffered saline and counterstained with TOPRO-iodide (Invitrogen) for 15 min. Cells were washed

three times in phosphate-buffered saline for 10 min; coverslips were added using VectaShield mounting medium (Vector Laboratories), and cells were examined by confocal microscopy using an Olympus FV-500 microscope with a  $\times$ 60 water objective. The experiment was repeated three times. Cells from at least two fields from each slide were counted by two individuals (one blind to treatment) and were scored as either negative or positive;  $\sim$ 200 cells per slide were counted. Cells were considered positive if multiple discrete puncta were observed. Negative cells were devoid of puncta, similar to cells incubated with normal rabbit serum.

**Analysis of Data**—The labeled bands from autoradiography were scanned using a GS-710 calibrated imaging densitometer, and the intensity of bands was quantitated by densitometric analysis using Quantity One software (Bio-Rad Laboratories). Equal exposure times for the autoradiograms were used when comparing the results of synthesis and protein secretion for LH, LH $\Delta$ T, FSH, and FSH-L dimers. The secretion  $t_{1/2}$  for the dimers corresponds to the time when 50% of the labeled dimer, as determined from the presence of  $\alpha$  subunit, accumulated in the medium when immunoprecipitating with CG $\beta$  antisera. Recovery of subunit (%) was determined as the amount of labeled subunit in medium as a fraction of the total. The sorting index of LH, LH $\Delta$ T, FSH, and FSH-L dimer corresponds to the ratio of the band intensity on autoradiograms obtained with forskolin (+F) to the band intensity obtained without (–F) forskolin. For immunofluorescence, results are expressed as the percentage of positive cells. Each experiment was repeated three to eight times, analyzed using a paired  $t$  test, and the results are expressed as mean  $\pm$  S.E., with  $p < 0.05$  considered significantly different.

## RESULTS

**Secretion of LH and LH $\Delta$ T After Steady-state Labeling**—Previously we showed in CHO cells that deleting the seven-amino acid hydrophobic carboxyl-terminal extension from the LH $\beta$  subunit enhances secretion and alters processing of the N-linked oligosaccharides of this truncated subunit (designated LH $\beta$  $\Delta$ T, Fig. 1) compared with the wild-type LH $\beta$  subunit (19). Thus, we suspected that, because this sequence affects the intracellular disposition of the subunit, it contributes to LH sorting in the regulated pathway. Because CHO cells lack a regulated secretory pathway, we examined the secretion of LH and the mutant LH $\Delta$ T in GH $_3$  cells, which contain both constitutive and regulated routes (20, 21). Fig. 2 shows the secretion of two sets of representative clones (*Groups 1* and *2*) expressing either LH or LH $\Delta$ T heterodimers after 16 h of labeling with [ $^{35}$ S]cysteine.

Immunoprecipitation of LH dimer secreting cells with  $\alpha$  antiserum co-precipitates the  $\beta$  subunit, reflecting the extent of heterodimer formation (Fig. 2, *A* and *B*, lane 2), and the combined and free (*asterisks*) forms of  $\alpha$  subunit. The unassembled  $\alpha$  subunit is unique in that it undergoes a post-translational modification resulting in a more heterogeneous form, which clearly distinguishes it from the dimer species (Fig. 2, *A* and *B*, lane 2). The free  $\alpha$  subunit is rapidly secreted with no detectable intracellular accumulation (Fig. 2, *A* and *B*, lane 1). As reported previously (18, 20), after overnight labeling the

## Sorting Signal for LH

secreted heterodimer precipitated with  $\alpha$  antiserum contains labeled  $\alpha$  subunit associated with only a fraction of the total labeled  $\beta$  subunit seen in the lysate (Fig. 2, A and B, compare lanes 2 and 3). This is presumably due to the presence of a stable, nonlabeled intracellular pool of assembly-competent LH $\beta$  subunit (see "Discussion"). Precipitation of LH dimer by  $\beta$  subunit antiserum co-precipitates  $\alpha$  subunit from the lysate and medium (Fig. 2, A and B, lanes 3 and 4). Note that the  $\alpha$  subunit is only weakly observed in the lysate (Fig. 2, A and B, lane 3), reflecting the extent of heterodimer accumulating after steady state labeling. Note also the absence of the heterogeneous free  $\alpha$  subunit in the media (Fig. 2, A and B, compare lanes 2 and 4). In contrast to the free  $\alpha$  subunit, a significant fraction of the unassembled  $\beta$  subunit accumulates intracellularly (Fig.

2, A and B, lane 3), despite the presence of excess  $\alpha$  subunit, and is confined to the ER (12, 14). This result agrees with earlier studies showing that in transfected cell lines the LH $\beta$  subunit inefficiently heterodimerizes with the  $\alpha$  subunit (18, 19) and the unassembled  $\beta$  subunit is not secreted efficiently.

As expected from our previous work in CHO cells (18), the extent of heterodimer formation is significantly increased when the heptapeptide is deleted (Fig. 2, A and B, lanes 5–8). This is reflected in the  $3.1 \pm 0.4$ -fold reduction ( $p < 0.05$ ) in the lysate fraction of the unassembled LH $\beta\Delta T$  subunit immunoprecipitated with  $\beta$  antiserum compared with the LH $\beta$  subunit (Fig. 2, A and B, lanes 3 and 7). In addition, immunoprecipitation with  $\alpha$  antiserum demonstrated a  $2.6 \pm 0.1$ -fold decrease ( $p < 0.05$ ) in the amount of intracellular LH $\Delta T$  (Fig. 2, A and B, lane 5) and  $2.3 \pm 0.2$ -fold increase ( $p < 0.05$ ) in the secretion of LH $\Delta T$  compared with LH dimer (lanes 2 and 6). This is associated with an increase in the incorporation of the labeled LH $\beta\Delta T$  subunit in heterodimer compared with the wild-type LH $\beta$  subunit. It is curious that the truncated LH $\beta\Delta T$  subunit migrates slower in the gel compared with the wild-type LH $\beta$  subunit. It is unclear whether this is due to the unmasking of a site for post-translational change or an effect on the mobility of the subunit on the gel. These data show that the total LH $\Delta T$  secreted from GH<sub>3</sub> cells is greater than LH dimer.

**Pulse-Chase Kinetics**—The steady-state labeling suggested that less of the LH $\Delta T$  dimer is retained in GH<sub>3</sub> cells consistent with an increased constitutive secretion. To assess this point more directly, we examined the kinetics of LH and LH $\Delta T$  secretion by pulse-chase labeling (Fig. 3). GH<sub>3</sub> cells were pulse-labeled for 20 min with [<sup>35</sup>S]cysteine and chased for the indicated times (Fig. 3). As expected in the case of wild-type LH (Fig. 3A), there is an excess accumulation of intracellular  $\beta$  subunit compared with the  $\alpha$  subunit. The  $t_{1/2}$  for secretion of LH was  $3.5 \pm 0.06$  h based on the appearance of dimer (determined from co-precipitated  $\alpha$  subunit) in the media. It is also evident that labeled  $\alpha$  subunit was chased into the secreted dimer more quickly than the labeled  $\beta$  subunit. Because  $\beta$  subunit antiserum was used, the labeled  $\alpha$  subunit detected at the earliest time point is likely combining with predominantly unlabeled assembly-competent LH $\beta$  subunit as described previously (15) (see also "Discussion"). In the case of LH $\Delta T$  (Fig. 3B), similar to earlier reports in which CHO cells were used, the efficiency of assembly was doubled as compared with LH (only 17% of the LH $\beta$  subunit assembled with the  $\alpha$  subunit versus 32% of LH $\beta\Delta T$ ), and its secretion into the media was faster ( $t_{1/2} = 1.8 \pm 0.21$  h) compared with LH. The recovery of both proteins in the media was greater than 90%, and thus differences in the secretion of LH and LH $\Delta T$  are not likely related to degradation. The labeled LH $\beta\Delta T$  subunit appeared




FIGURE 1. Schematic diagram of the gonadotropin subunits. These include:  $\alpha$  subunit; LH $\beta$ , luteinizing hormone  $\beta$  subunit (the cross-hatched area of region 115–121 denotes the hydrophobic heptapeptide); LH $\beta\Delta T$ , carboxyl-terminal mutant of LH $\beta$  subunit truncated at amino acid 114; FSH $\beta$ , follicle-stimulating hormone  $\beta$  subunit; FSH $\beta$ -L, the cross-hatched area of region 112–118 denotes the hydrophobic heptapeptide of LH $\beta$ . N, Asn-linked oligosaccharides.





FIGURE 2. Secretion of representative clones (Group 1 (A) and Group 2 (B)) expressing either LH ( $n = 2$  clones) or LH $\Delta T$  ( $n = 2$  clones) dimers after being labeled for 16 h with  $20 \mu\text{Ci/ml}$  [<sup>35</sup>S]cysteine. Media (M) and lysates (L) were immunoprecipitated with anti- $\alpha$  and CG $\beta$  sera. The immunoprecipitates were resolved by SDS-PAGE and subjected to autoradiography. The migration of  $\alpha$  and  $\beta$  subunits and the molecular mass markers are indicated. The data shown are representative of five independent experiments. Asterisks represent free  $\alpha$  subunit.

earlier in the secreted dimer as compared with wild-type  $\beta$  subunit in the LH dimer. Taken together, the data suggest that removal of the heptapeptide increased secretion through the constitutive pathway and that this sequence is associated with directing LH to the regulated pathway.


**Secretagogue-stimulated Secretion of LH/LH $\Delta$ T**—Earlier we showed that LH accumulates in a secretagogue-sensitive pool, reflecting segregation to the regulated pathway (20, 21). If deleting the heptapeptide leads to increased constitutive secretion, as proposed above, the amount of forskolin-releasable LH $\Delta$ T dimer should be decreased compared with LH. To investigate this issue we labeled the cells with [ $^{35}$ S]sulfate (Fig. 4). Because the LH N-linked oligosaccharides are sulfated in GH $_3$  cells (20), we reasoned that, given that this modification occurs in the trans-Golgi and is a final step in the maturation of the oligosaccharides just prior to secretion, only those mature

forms of LH/LH $\Delta$ T accumulating in the late stages of the secretion pathway will contain [ $^{35}$ S]sulfate. Presumably this enriched population is in a secretagogue-releasable pool, and as a result the background would be reduced compared with [ $^{35}$ S]cysteine labeling. After a 16-h labeling, media and lysates collected from one well of each dimer served as the 0 time controls for the subsequent experiments. The distribution of the intracellular and secreted proteins is comparable with that seen for the cysteine labeled products (see Fig. 2). To reduce the background of constitutive secretion after labeling, LH and LH $\Delta$ T cells were preincubated for 2 h in chase medium. At the end of the initial 2 h of preincubation, prior to forskolin addition, cells from the control well and the well to receive forskolin secreted comparable amounts of either LH or LH $\Delta$ T into the medium (Fig. 4, A and B, lanes 1 and 2, asterisks). The addition of forskolin during the second 2-h period stimulated release of sulfate-labeled LH dimer ( $2.9 \pm 0.4$ -fold,  $p < 0.05$ ) compared with cells incubated without forskolin (Fig. 4A, lanes 3 and 4; Table 1). Importantly, the increased secretion was accompanied by a decrease of intracellular LH ( $3.2 \pm 0.2$ -fold,  $p < 0.05$ ) compared with un-stimulated cells (Fig. 4A, lanes 5 and 6; Table 1). It is also noteworthy that no shift in electrophoretic migration between the intracellular and secreted subunits was detected, indicating that both intracellular forms were fully processed to complex oligosaccharides, and thus the mature proteins accumulated intracellularly after traversing the late Golgi compartment (20).

In contrast to LH dimer, there was minimal stimulation of LH $\Delta$ T secretion by forskolin ( $1.25 \pm 0.07$ -fold versus no forskolin; Fig. 4B, lanes 3 and 4; Table 1). In addition, the unstimulated pool for LH $\Delta$ T during the forskolin stimulation period was reduced by  $4.4 \pm 0.5$ -fold ( $p < 0.05$ ) compared with LH dimer (Fig. 4, B, lane 5, versus A, lane 5). Although it is clear that the forskolin-stimulated release of LH $\Delta$ T was reduced, it was difficult to accurately quantitate the density of bands (even after the prolonged exposure) because of its decreased intracellular accumulation during the entire 4-h incubation. To address this issue, an additional experiment was performed where the initial 2-h preincubation period was eliminated. Cells expressing LH $\Delta$ T were incubated for only 2 h in the presence or absence of forskolin immediately after a 16-h labeling (Fig. 4C). In this case, there was a larger pool of intracellular LH $\Delta$ T, compared with the 4-h chase, as reflected by the increase in the nonstimulated level of LH $\Delta$ T (Fig. 4C, lane 3). However, in the presence of forskolin, the secretion of LH $\Delta$ T was stimulated only  $1.39 \pm 0.11$ -fold compared with nonstimulated cells (Fig. 4C, lanes 1 and 2) and accompanied by a  $1.38 \pm 0.14$ -fold decrease in the



**FIGURE 3. Pulse-chase kinetics of LH (A) and LH $\Delta$ T (B) dimers.** GH $_3$  cells were pulse-labeled with  $80 \mu\text{Ci/ml}$  [ $^{35}$ S]cysteine for 20 min and then chased for the indicated times. Chase at 0 h indicates the lysate sample prepared immediately after the pulse. Lysates and media were immunoprecipitated with CG $\beta$  antiserum and subjected to SDS-PAGE followed by autoradiography. The migration of  $\alpha$  and  $\beta$  subunits is indicated. The  $t_{1/2}$  was calculated from the appearance of the  $\alpha$  subunit in the media, which indicates the extent of dimer formation. The data shown are representative of three independent experiments.



**FIGURE 4. Secretagogue-stimulated secretion of LH and LH $\Delta$ T dimers from GH $_3$  cells.** The top part of the figure illustrates the experimental protocol. Cells were incubated for 16 h with [ $^{35}$ S]sulfate ( $0.7 \text{ mCi/ml}$ ) followed by a 2-h incubation in chase media. Media (M, asterisks) of LH (A) and LH $\Delta$ T (B) collected at the end of the 2-h preincubation period demonstrate that cells in each pair of wells synthesized the same amount of LH/LH $\Delta$ T prior to forskolin addition. Subsequently, these media were replaced with fresh chase media, and the incubation was extended for an additional 2 h in the presence (+, lanes 4 and 6) or absence (–, lanes 3 and 5) of  $25 \mu\text{M}$  forskolin. Panel C represents LH $\Delta$ T incubated in the presence (+, lanes 2 and 4) or absence (–, lanes 1 and 2) of forskolin for 2 h without preincubation. All cell lysates (L) and media (M) were immunoprecipitated with  $\alpha$  antiserum. The data shown are representative of four independent experiments.

## Sorting Signal for LH

**TABLE 1**

**Forskolin (F)-stimulated secretion of dimers from transfected GH<sub>3</sub> cells**


<sup>35</sup>S-Sulfated (LH, LHΔT) or cysteine-labeled (FSH, FSH-L) and immunoprecipitated proteins from x-rays similar to those shown in Figs. 5, A and B, and 8B, respectively, were quantitated by densitometry. The sorting index (+F/−F) values for medium correspond to the ratio of the band intensity (volume OD/mm<sup>2</sup>) obtained with forskolin-stimulated release (+F; 25 μM) to the band intensity obtained without forskolin (−F). Values for lysates (−F/+F) represent the decrease of accumulation of protein after forskolin stimulation. The data represent the mean values of four independent experiments ± S.E. \*, *p* < 0.05 for LH vs. LHΔT and FSH vs. FSH-L.

Protein	Sorting index	
	Medium (+F/−F)	Lysate (−F/+F)
LH	2.90 ± 0.40*	3.20 ± 0.23*
LHΔT	1.20 ± 0.07	1.15 ± 0.06
FSH	1.05 ± 0.04	1.24 ± 0.05
FSH-L	2.24 ± 0.13*	2.09 ± 0.12*


lysate pool (lanes 3 and 4). These results confirm the 4-h chase data for LHΔT showing that the intracellular stores of LHΔT are significantly reduced relative to LH dimer. Similar differences were observed after forskolin-stimulated secretion with cysteine-labeled LH and LHΔT dimers (data not shown). Overall, these experiments indicate that the absence of the heptapeptide in the LHβ subunit leads to less regulated secretion of LHΔT.

**Immunofluorescence**—We next examined whether the constitutive nature of LHΔT described above could be observed using immunofluorescence. Before fixation, cells were incubated for 4 h in the presence or absence of CHX to inhibit protein synthesis. This experiment predicts that in the absence of newly synthesized protein, the extent of LHΔT secretion will be greater than LH, thus having less intracellular retention. Untreated GH<sub>3</sub> cells expressing LH or LHΔT have a similar fluorescence pattern with randomly dispersed puncta (Fig. 5, A and D). These puncta presumably represent post-Golgi LH- or LHΔT-containing vesicles and are not observed in cells incubated with normal rabbit serum instead of primary antiserum (Fig. 5, C and F). Note that for Fig. 5, A, B, D, and E, the counterstaining of the nucleus is not shown to better visualize the green puncta. The percentage of positive cells is similar between untreated and 4-h CHX-treated cells (82.4 ± 4.5 versus 77.3 ± 2.1%). In contrast, the percentage of LHΔT positive cells decreases from 80.4 ± 5.2% in untreated cells to 57.3 ± 4.5% after 4 h of CHX treatment (*p* < 0.05; Fig. 6), reflecting less regulated secretion of LHΔT.

**Forskolin-stimulated Secretion of FSH/FSH-L**—We next asked whether the LHβ heptapeptide was sufficient to reroute a constitutively secreted protein to the regulated pathway. To examine this question, we fused the LHβ heptapeptide to the carboxyl terminus of the FSHβ subunit (FSHβ-L; Fig. 1). In this experiment, two parameters were compared between FSH-L and wild-type FSH: the extent of intracellular accumulation and the sensitivity to forskolin stimulation. Previously we showed that after stimulation with forskolin, a small fraction of newly synthesized FSH was secretagogue-sensitive and thus entered the regulated pathway (21). Those analyses used polyclonal antisera to FSH/FSHβ. Here we used a high affinity FSH dimer-specific monoclonal antibody (23, 24) to identify the proteins because it does not cross-react with the uncombined FSHβ subunit, thus reducing the background. As shown previously,




**FIGURE 5. Immunofluorescence of untreated and 4-h CHX (15 μg/ml)-treated LH and LHΔT GH<sub>3</sub> cells.** Untreated LH (A) and LHΔT (D) cells have a similar fluorescence pattern with randomly dispersed puncta (green) around the nucleus. Insets show a higher magnification of LH and LHΔT cells for greater definition of the puncta. Counterstaining of the nucleus (red; C and F) is not shown in A, B, D, and E to enhance the contrast of the puncta. These puncta were not observed in cells incubated with normal rabbit serum (NRS; C and F) instead of primary antibody.




**FIGURE 6. Percentage of LH- or LHΔT-positive cells after incubation in the presence or absence (Untreated) of CHX.** Results are the mean ± S.E. \*, *p* < 0.05.

the α and FSHβ subunits migrate at similar rates on SDS gels, and their resolution is not clear (21). It is evident that after 4 h of labeling and subsequent chase, more FSH-L accumulates in the lysate compared with FSH (Fig. 7B, lanes 3 and 7); this is accompanied by a corresponding decrease in the secretion of FSH-L (lane 5). Incubation with forskolin stimulated FSH-L 2.24 ±

### A. Experimental Protocol



### B. FSH 54B mAb



**FIGURE 7. Secretagogue-stimulated secretion of FSH and FSH-L dimers from GH<sub>3</sub> cells.** *A*, experimental protocol. *B*, represents FSH and FSH-L incubated in the presence (+, lanes 2, 4, 6, and 8) or absence (-, lanes 1, 3, 5, and 7) of forskolin. All cell lysates (L) and media (M) were immunoprecipitated with FSH 54B monoclonal antibody (mAb). Note that the  $\alpha$  and FSH $\beta$  subunits co-migrate on SDS gels at ~23 kDa. The data shown are representative of four independent experiments.

0.13-fold (Fig. 7*B*, lanes 5 and 6) and FSH 1.05  $\pm$  0.04-fold (lanes 1 and 2; Table 1). Moreover, there was a greater depletion of the intracellular pool (lysate) of FSH-L (2.09  $\pm$  0.12-fold; Fig. 7*B*, lanes 7 and 8) than FSH (1.24  $\pm$  0.05-fold; lanes 3 and 4) following forskolin stimulation (Table 1). Similar results were observed using different clones for FSH and FSH-L. These results indicate that the carboxyl-terminal seven amino acids of LH $\beta$  can reroute a fraction of FSH to the regulated pathway.

### DISCUSSION

An important functional feature of the gonadotropins is their characteristic secretion patterns. LH is released in pulses via the regulated pathway in response to the secretagogue GnRH, whereas FSH is secreted constitutively. Clearly, this routing is essential for the coordinated roles of LH and FSH in the reproductive tract. Defining the sorting signals governing this trafficking is critical for understanding the link between their secretion and reproductive function. Although no common linear amino acid sequences capable of diverting secretory proteins into the regulated secretory pathway have been identified, signaling information has been observed in numerous domains of several secretory proteins (25–32). Here we describe a novel sorting signal (comprising a carboxyl-terminal seven-amino acid peptide in the LH $\beta$  subunit) that contributes to the diver-

sion of newly synthesized LH to the regulated pathway of GH<sub>3</sub> cells.

How the heptapeptide functions as a sorting signal is not clear. It could act directly as a sorting signal or indirectly through changes in the conformation of the molecule. Although the latter option cannot be excluded, the ability of the heptapeptide to reroute a portion of FSH to the regulated pathway suggests that the sequence *per se* is a sorting determinant. Recently, it was shown that  $\alpha$ -helical sequences bearing stretches of hydrophobic amino acids at carboxyl-end regions can direct a variety of secretory proteins into the regulated secretion pathway (33). We examined the predicted secondary structure for the LH $\beta$  subunit with PSIPRED (protein structure prediction server) (34, 35). According to this analysis, the last 40 amino acids prefer a coil configuration. As such, it appears that the proposed sorting activity of the heptapeptide does not require a helical structure, although analogous to the above sorting sequence, it contains a hydrophobic cluster. Inspection of the heptapeptide sequence reveals a similarity to the sorting sequence XXXLL found in the cytoplasmic domain of numerous transmembrane proteins (36). Although recognition of the LH heptapeptide occurs in the luminal compartment rather than the cytoplasm, it is intriguing to consider that the heptapeptide has analogous recognition properties. For example, the heptapeptide might represent a granule-targeting domain that associates with a receptor in the vesicle membrane. It is likely that the heptapeptide works in concert with other signals in the LH molecule, because a portion of LH $\Delta$ T still enters the regulated pathway. Importantly, the sequence studied here is well conserved in mammals (37), consistent with a proposed sorting function.

Further supporting the sorting feature associated with the heptapeptide are the observations that although the DNA sequence of the LH $\beta$  subunit gene encodes the heptapeptide, this hydrophobic stretch is not detected in the intact dimer isolated from pituitary tissue or urine (38, 39). Moreover, in the latter study (39), heterogeneity among carboxyl-terminal residues 114–121 of the LH $\beta$  subunit has been observed. That absence of the heptapeptide leads to less regulated secretion, with a concomitant enhanced constitutive secretion, supports its role in governing the trafficking of the LH $\beta$  subunit in the gonadotrope.

One critical finding regarding the expression of the gonadotropin family from transfected cell lines is that, although LH $\beta$  and FSH $\beta$  are retained intracellularly as monomers, only the LH $\beta$  subunit is slow to assemble with the  $\alpha$  subunit (13, 15). Also, like the LH $\beta$  subunit, the presence of the heptapeptide on FSH $\beta$  decreases the extent of assembly with the  $\alpha$  subunit (data not shown). The other  $\beta$  subunits assemble with the  $\alpha$  subunit at an efficiency of more than 90%. Moreover, in transfected GH<sub>3</sub> cells expressing LH dimer, a significant fraction of the pulse-labeled  $\alpha$  subunit is secreted in association with unlabeled LH $\beta$  subunit, whereas the labeled  $\beta$  subunit remains unassociated. One explanation for the delay is the under-representation of labeled subunits by a large pool of nonlabeled species. Such pools of unassembled subunits have been observed in other systems (40–42). However, what is unusual in cells that contain LH dimers is that the population of free



LH $\beta$  subunit persists despite the presence of excess  $\alpha$  subunits. Thus, one explanation is that the LH $\beta$  subunit accumulates in at least two different luminal pools. Consistent with this view, several reports suggest that the ER is functionally mosaic and comprises specialized subdomains (43–48). These subcompartments may have distinct functions and a distinct distribution of resident and transient proteins. Thus, one model for how cells segregate LH/FSH heterodimers in the gonadotrope is assembly in distinct sub-ER compartments as proposed previously (23). This predicts that LH devoid of the heptapeptide will lose specificity in these domains. Alternatively, the selective routing of LH/FSH to the regulated/constitutive secretory pathway may be determined at the level of the trans-Golgi (49).

One issue that arises is whether *in vivo* LH and FSH dimers are routed via monohormone-containing vesicles or co-localize with the same carriers that bifurcate into monohormonal units. Given that our present experiments were performed with clonal GH $_3$  cells, gonadotrope heterogeneity is not a variable, implying that the existence of only monohormonal LH and FSH secretion is governed primarily by general regulated and constitutive release mechanisms, respectively. Therefore, if a substantial population of vesicles initially contain both hormones, then encoded motifs are likely responsible for dictating the recognition to their respective pathways. Further characterization of the cognate recognition molecules for sorting determinants will be essential for understanding the underlying mechanisms by which LH and FSH are differentially sorted. Such investigations will also provide more general insight into how the entry of proteins into regulated secretory compartments is controlled.

*Acknowledgments*—We thank Drs. Phyllis Hanson, T. Rajendra Kumar, and Rick Sifers for helpful comments. We are also grateful to Dennis Oakley for help with confocal microscopy and to Laura Kyro, Barbara Machalek, Amanda Fisher, and Linda Lobos for preparation of the manuscript.

## REFERENCES

- Pierce, J. G., and Parsons, T. F. (1981) *Annu. Rev. Biochem.* **50**, 465–495
- Sairam, M. R. (1983) in *Hormonal Proteins and Peptides: Gonadotropic Hormones* (Li, C. H., ed) pp. 1–79, Academic Press, New York
- Green, E. D., Boime, I., and Baenziger, J. U. (1986) *Mol. Cell. Biochem.* **72**, 81–100
- Kessler, M. J., Mise, T., Ghai, R. D., and Bahl, O. P. (1979) *J. Biol. Chem.* **254**, 7909–7914
- Parsons, T. F., and Pierce, J. G. (1980) *Proc. Natl. Acad. Sci. U. S. A.* **77**, 7089–7093
- Luderer, U., and Schwartz, N. B. (1992) in *Serono Symposium on FSH: Regulation of Secretion and Molecular Mechanisms of Action* (Hunzicker-Dunn, M., and Schwartz, N. B., eds) pp. 1–25, Springer-Verlag, New York
- Farnworth, P. G. (1995) *J. Endocrinol.* **145**, 387–395
- Currie, R. J., and McNeilly, A. S. (1995) *J. Endocrinol.* **147**, 259–270
- Thomas, S. G., and Clarke, I. J. (1997) *Endocrinology* **138**, 1347–1350
- Baenziger, J. U., Kumar, S., Brodbeck, R. M., Smith, P. L., and Beranek, M. C. (1992) *Proc. Natl. Acad. Sci. U. S. A.* **89**, 334–338
- Baenziger, J. U. (1994) *Faseb. J.* **8**, 1019–1025
- Talmdage, K., Vamvakopoulos, N. C., and Fiddes, J. C. (1984) *Nature* **307**, 37–40
- Matzuk, M. M., Spangler, M. M., Camel, M., Suganuma, N., and Boime, I. (1989) *J. Cell Biol.* **109**, 1429–1438
- Hayashizaki, Y., Miyai, K., Kato, K., and Matsubara, K. (1985) *FEBS Lett.* **188**, 394–400
- Corless, C. L., Matzuk, M. M., Ramabhadran, T. V., Krichevsky, A., and Boime, I. (1987) *J. Cell Biol.* **104**, 1173–1181
- Keene, J. L., Matzuk, M. M., Otani, T., Fauser, B. C., Galway, A. B., Hsueh, A. J., and Boime, I. (1989) *J. Biol. Chem.* **264**, 4769–4775
- Kaetzel, D. M., Virgin, J. B., Clay, C. M., and Nilson, J. H. (1989) *Mol. Endocrinol.* **3**, 1765–1774
- Muyan, M., Furuhashi, M., Sugahara, T., and Boime, I. (1996) *Mol. Endocrinol.* **10**, 1678–1687
- Matzuk, M. M., Krieger, M., Corless, C. L., and Boime, I. (1987) *Proc. Natl. Acad. Sci. U. S. A.* **84**, 6354–6358
- Bielinska, M., Rzymkiewicz, D., and Boime, I. (1994) *Mol. Endocrinol.* **8**, 919–928
- Muyan, M., Rzymkiewicz, D. M., and Boime, I. (1994) *Mol. Endocrinol.* **8**, 1789–1797
- Dannies, P. S. (1999) *Endocr. Rev.* **20**, 3–21
- Garcia-Campayo, V., Jablonka-Shariff, A., and Boime, I. (2004) *J. Biol. Chem.* **279**, 44286–44293
- Jablonka-Shariff, A., Kumar, T. R., Eklund, J., Comstock, A., and Boime, I. (2006) *Mol. Endocrinol.* **20**, 1437–1446
- Childs, G. R. (1985) in *Endocrinology* (Labrie, F., and Proulx, L., eds) pp. 499–503, Elsevier, Amsterdam
- Arvan, P., Zhang, B. Y., Feng, L., Liu, M., and Kuliawat, R. (2002) *Curr. Opin. Cell Biol.* **14**, 448–453
- Tooze, S. A., Martens, G. J., and Huttner, W. B. (2001) *Trends Cell Biol.* **11**, 116–122
- Dannies, P. S. (2001) *Mol. Cell. Endocrinol.* **177**, 87–93
- Cool, D. R., Normant, E., Shen, F., Chen, H. C., Pannell, L., Zhang, Y., and Loh, Y. P. (1997) *Cell* **88**, 73–83
- Brechler, V., Chu, W. N., Baxter, J. D., Thibault, G., and Reudelhuber, T. L. (1996) *J. Biol. Chem.* **271**, 20636–20640
- Bundgaard, J. R., Birkeedal, H., and Rehfeld, J. F. (2004) *J. Biol. Chem.* **279**, 5488–5493
- Assadi, M., Sharpe, J. C., Snell, C., and Loh, Y. P. (2004) *Biochemistry* **43**, 7798–7807
- Dikeakos, J. D., Lacombe, M. J., Mercure, C., Mireuta, M., and Reudelhuber, T. L. (2007) *J. Biol. Chem.* **282**, 1136–1143
- McGuffin, L. J., Bryson, K., and Jones, D. T. (2000) *Bioinformatics (Oxf)* **16**, 404–405
- Jones, D. T. (1999) *J. Mol. Biol.* **292**, 195–202
- Bonifacino, J. S., and Traub, L. M. (2003) *Annu. Rev. Biochem.* **72**, 395–447
- Watanabe, N., Hatano, J., Asahina, K., Iwasaki, T., and Hayakawa, S. (2007) *Comp. Biochem. Physiol. A Mol. Integr. Physiol.* **146**, 105–118
- Keutmann, H. T., Williams, R. M., and Ryan, R. J. (1979) *Biochem. Biophys. Res. Commun.* **90**, 842–848
- Pantel, J., Robert, P., Troalen, F., Kujas, M., Bellet, D., and Bidart, J. M. (1998) *Endocrinology* **139**, 527–533
- Tartakoff, A., and Vassalli, P. (1979) *J. Cell Biol.* **83**, 284–299
- Ho, M. K., and Springer, T. A. (1983) *J. Biol. Chem.* **258**, 2766–2769
- Kvist, S., Wiman, K., Claesson, L., Peterson, P. A., and Dobberstein, B. (1982) *Cell* **29**, 61–69
- Sitia, R., and Meldolesi, J. (1992) *Mol. Biol. Cell* **3**, 1067–1072
- Griffiths, G. (1996) *Protoplasma* **195**, 37–58
- Cabral, C. M., Choudhury, P., Liu, Y., and Sifers, R. N. (2000) *J. Biol. Chem.* **275**, 25015–25022
- Kamhi-Nesher, S., Shenkman, M., Tolchinsky, S., Fromm, S. V., Ehrlich, R., and Lederkremer, G. Z. (2001) *Mol. Biol. Cell* **12**, 1711–1723
- Shimizu, Y., and Hendershot, L. M. (2007) *Adv. Exp. Med. Biol.* **594**, 37–46
- Avezov, E., Frenkel, Z., Ehrlich, M., Herscovics, A., and Lederkremer, G. Z. (2008) *Mol. Biol. Cell* **19**, 216–225
- Nakatsu, F., and Ohno, H. (2003) *Cell Struct. Funct.* **28**, 419–429